

Main Examination period 2019

## MTH5123: Differential Equations

**Duration: 2 hours**

Student number

--	--	--	--	--	--	--	--	--

Desk number

--	--	--

---

**Apart from this page, you are not permitted to read the contents of this question paper until instructed to do so by an invigilator.**

**Write your solutions in the spaces provided in this exam paper. If you need more paper, ask an invigilator for an additional booklet and attach it to this paper at the end of the exam.**

<p><b>You should attempt ALL questions. Marks available are shown next to the questions.</b></p>
--

**Calculators are not permitted in this examination. The unauthorised use of a calculator constitutes an examination offence.**

Complete all rough work in the answer book and cross through any work that is not to be assessed.

Possession of unauthorised material at any time when under examination conditions is an assessment offence and can lead to expulsion from QMUL. Check now to ensure you do not have any unauthorised notes, mobile phones, smartwatches or unauthorised electronic devices on your person. If you do, raise your hand and give them to an invigilator immediately.

It is also an offence to have any writing of any kind on your person, including on your body. If you are found to have hidden unauthorised material elsewhere, including toilets and cloakrooms, it will be treated as being found in your possession. Unauthorised material found on your mobile phone or other electronic device will be considered the same as being in possession of paper notes. A mobile phone that causes a disruption in the exam is also an assessment offence.

**Exam papers must not be removed from the examination room.**

**Examiners: Shabnam Beheshti, Weini Huang**

---

**This page is for marking purposes only.  
Do not write on it.**

<b>Question</b>	<b>Mark</b>	<b>Comments</b>
<b>1</b>	<b>/ 20</b>	
<b>2</b>	<b>/ 20</b>	
<b>3</b>	<b>/ 20</b>	
<b>4</b>	<b>/ 20</b>	
<b>5</b>	<b>/ 20</b>	
<b>Total</b>		

**Note for the rest of this exam paper ODE refers to ordinary differential equation.**

**Question 1. [20 marks]** The equation of motion for a falling object of mass  $m$  is given by

$$m \frac{dv}{dt} = mg - \gamma v,$$

where  $g = 9.8 \text{ m/s}^2$  is the acceleration due to gravity,  $\gamma$  is a constant called the drag coefficient and  $v = v(t)$  denotes the velocity of the object at time  $t$ . Assume  $m = 10 \text{ kg}$  and  $\gamma = 2 \text{ kg/s}$ .

- (a) Find the general solution to this differential equation with the given constants. [5]
- (b) Now find the specific solution satisfying the initial condition  $v(0) = 49$ . [5]
- (c) Draw integral curves in the  $t$ - $v$  plane for various initial conditions, including the initial condition  $v(0) = 49$ . [5]
- (d) Interpret your graph for this model, explaining briefly the behaviour of your solutions as  $t \rightarrow \infty$  for different initial conditions. [5]

**Write your solutions here**

**Continue-1: solutions to question 1**

**Write your solutions here**

**Continue-2: solutions to question 1**

**Write your solutions here**

**Question 2. [20 marks]** Consider the following first-order, linear, inhomogeneous initial value problem

$$\begin{cases} y' &= y \tan x + \sin x, & -\pi/2 < x < \pi/2 \\ y(0) &= 1 \end{cases} .$$

- (a) Find a solution  $y = y(x)$  to the initial value problem. [10]
- (b) Use the Picard-Lindelöf Theorem to justify existence and uniqueness of solutions to the above IVP in an appropriate rectangular domain. [10]

**Write your solutions here**

**Continue-1: solutions to question 2**

**Write your solutions here**

**Continue-2: solutions to question 2**

**Write your solutions here**



**Continue-3: solutions to question 2**

**Write your solutions here**

**Question 3. [20 marks]** Consider the differential equation given by

$$x^2 + \frac{f(y)}{xy} + \ln |xy| \frac{dy}{dx} = 0.$$

- (a) Find all functions  $f(y)$  such that the differential equation becomes exact. [10]
- (b) For the function  $f$  which makes the differential equation exact and which further satisfies  $f(1) = 1$ , solve the equation in implicit form. [10]

**Write your solutions here**

**Continue-1: solutions to question 3**

**Write your solutions here**

**Continue-2: solutions to question 3**

**Write your solutions here**

**Question 4. [20 marks]** Consider the following Euler-type equation

$$x^2 \frac{d^2 y}{dx^2} - 2y = 0. \quad (*)$$

(a) Using  $x = e^t$  and  $z(t) = y(e^t)$ , verify that (\*) can be rewritten as

$$\ddot{z} - \dot{z} - 2z = 0.$$

Use this equation to find the general solution  $y = y(x)$  to (\*). [5]

(b) Next, consider the Boundary Value Problem (BVP) for the second order inhomogeneous differential equation

$$x^2 \frac{d^2 y}{dx^2} - 2y = f(x), \quad y(1) = 0, \quad y(2) + 2y'(2) = 0.$$

Formulate the corresponding left-end and right-end initial value problems. You **do not** need to solve the IVPs. [5]

(c) Assume that

$$y_L(x) = \frac{1}{3x} - \frac{1}{3}x^2 \quad \text{and} \quad y_R(x) = \frac{4}{x}$$

are the solutions to the IVPs in part (b). Write down the Green's function  $G(x, s)$  for the BVP in simplified form. [5]

(d) Represent the solution to the BVP in terms of the Green's function  $G(x, s)$  for the particular choice  $f(x) = e^x$ . You **do not** need to evaluate the resulting integrals. [5]

**Write your solutions here**

**Continue-1: solutions to question 4**

**Write your solutions here**

**Continue-2: solutions to question 4**

**Write your solutions here**

**Continue-3: solutions to question 4**

**Write your solutions here**



**Continue-4: solutions to question 4**

**Write your solutions here**

**Question 5. [20 marks]** Consider the autonomous dynamical system given by

$$\dot{x} = 4y, \quad \dot{y} = -x.$$

- (a) Rewrite the system in matrix form and find the associated eigenvalues and eigenvectors. [5]
- (b) Determine the solutions of the corresponding initial value problems for the general initial conditions  $x(0) = a, y(0) = b$ . [5]
- (c) Sketch the phase portrait in the  $(x, y)$  phase plane and describe the shape of the trajectories in the phase plane. If the initial condition is  $(x(0), y(0)) = (3, 4)$ , describe the qualitative behaviour of the solution given in the phase portrait. [5]
- (d) Determine all fixed points of the system and describe the stability of  $(x(t), y(t)) = (0, 0)$  as a solution for the linear system. What type of equilibrium point is  $(0, 0)$ ? [5]

**Write your solutions here**

**Continue-1: solutions to question 5**

**Write your solutions here**

**Continue-2: solutions to question 5**

**Write your solutions here**

---

**End of Paper – An appendix of 1 page follows.**

## Appendix

**Picard-Lindelöf Theorem.** Let  $\mathcal{D}$  be the rectangular domain in the  $xy$  plane defined as  $\mathcal{D} = (|x - a| \leq A, |y - b| \leq B)$  and suppose  $f(x, y)$  is a function defined on  $\mathcal{D}$  which satisfies the following conditions:

- (i)  $f(x, y)$  is continuous and therefore bounded in  $\mathcal{D}$
- (ii) the parameters  $A$  and  $B$  satisfy  $A \leq B/M$  where  $M = \max_{\mathcal{D}} |f(x, y)|$
- (iii)  $|\frac{\partial f}{\partial y}|$  is bounded in  $\mathcal{D}$ .

Then there exists a unique solution on  $\mathcal{D}$  to the initial value problem

$$\frac{dy}{dx} = f(x, y), \quad y(a) = b.$$

**Green's Function Method.** If there exists a unique solution  $y(x)$  to the inhomogeneous **boundary value problem**

$$\mathcal{L}(y) = a_2(x)y'' + a_1(x)y' + a_0(x)y = f(x)$$

in an interval  $x \in [x_1, x_2]$  with linear homogeneous boundary conditions

$$\alpha y'(x_1) + \beta y(x_1) = 0, \quad \gamma y'(x_2) + \delta y(x_2) = 0,$$

it can be found by the Green's function method:

$$y(x) = \int_{x_1}^{x_2} G(x, s) f(s) ds, \quad G(x, s) \equiv \begin{cases} A(s) y_L(x), & x_1 \leq x \leq s \\ B(s) y_R(x), & s \leq x \leq x_2 \end{cases}.$$

Here

$$A(s) = \frac{y_R(s)}{a_2(s)W(s)}, \quad B(s) = \frac{y_L(s)}{a_2(s)W(s)}, \quad W(s) \equiv y_L(s)y'_R(s) - y_R(s)y'_L(s)$$

and  $y_L(x), y_R(x)$  are solutions to the left/right initial value problems:

$$\mathcal{L}(y) = 0, \quad y(x_1) = \alpha, \quad y'(x_1) = -\beta; \quad \text{and} \quad \mathcal{L}(y) = 0, \quad y(x_2) = \gamma, \quad y'(x_2) = -\delta.$$

**End of Appendix.**