

## LECTURE 9

### 5 Conditional Heteroscedastic Models

The models referred to as conditional heteroscedastic models are used for modeling the volatility of asset returns. The volatility means the conditional standard deviation of the underlying asset return. It plays important role in

- calculating value at risk
- asset allocation
- helps to improve the efficiency in parameter estimation and the accuracy in interval forecast.

The volatility index VIX is a financial instrument traded in financial markets.



— ViX represents markets expectations for volatility over the coming 30 days; VIX is used to measure the level of risk or stress in the market when making investment decisions.


We shall discuss:


- the autoregressive conditional heteroscedastic model (ARCH) of Engle (1982)
- the generalized ARCH model (GARCH) by Bollerslev (1982) and show some applications.

## 5.1 Characteristics of Volatility

The feature of volatility is that volatility is not directly observable. It has some characteristics that are commonly seen in asset returns:

- There exist volatility clusters (volatility might be high for certain time periods and low for other periods)  
"large changes tend to be followed by large changes, of either sign, and small changes tend to be followed by small changes" Mandelbrot
- Volatility evolves in continuous manner, volatility jumps are rare.
- Volatility does not diverge to infinity, it remains within some fixed range. Statistically speaking, volatility is often stationary.
- It reacts differently to a big price increase or a big price drop. This effect is called leverage effect.





S&P 500 VOLATILITY CLUSTERING 90 ...  
researchgate.net

These properties play an important role in developing volatility models, which are trying to capture mentioned characteristics.

### 5.1.1 Structure of the models

Let  $r_t$  be the log return of an asset  $P_t$  at time  $t$ :

$$r_t = \log P_t - \log P_{t-1}$$

Basic idea behind volatility study:

- series of log returns  $r_t$  is uncorrelated or with minor lower order correlation,
- but series of log returns  $r_t$  is dependent series.

**Example.** Figure 3.4 shows the ACF of the monthly stock returns of Intel corporation for 1973 to 2003:

- Figure 3.4 a) shows the sample ACF of the returns  $r_t$  which suggests no significant correlation
- Figure 3.4 c) shows the sample ACF of the absolute log return  $|r_t|$  which is correlated
- Figure 3.4 b) shows the sample ACF of the squared returns  $r_t^2$  which suggest correlation


**Conclusion:** Figures show that monthly returns  $r_t$  are serially uncorrelated, but dependent. Volatility models try to capture such dependence.

**Notice:** if  $e_t$  are independent variables then

$e_t^2, |e_t|$  are also independent variables.

Independent variables are uncorrelated. Therefore all three series

$e_t, e_t^2, |e_t|$  are uncorrelated.



**Figure 3.4.** Model checking statistics of the Gaussian ARCH(1) model in Eq. (3.11) for the monthly log returns of Intel stock from January 1973 to December 2003: parts (a), (b), and (c) show the sample ACF of the standardized residuals, their squared series, and absolute series, respectively; part (d) is the time plot of standardized residuals.






Example. Plot of daily log returns  $r_t$  of S&P index

Sample ACF of  $r_t$  does not show correlation in  $r_t$ .

Sample ACF of  $|r_t|$  shows strong correlation in  $|r_t|$ .

Sample ACF of  $r_t^2$  would also show strong correlation in  $r_t^2$ .



## Definition of volatility.

We introduce

- conditional mean  $\mu_t$
- conditional variance  $\sigma_t^2$

given the information  $F_{t-1}$  available at time  $t - 1$ :

$$\mu_t = E(r_t | F_{t-1}), \quad \sigma_t^2 = \text{Var}(r_t | F_{t-1}) = E[(r_t - \mu_t)^2 | F_{t-1}].$$

We observed in previous example that serial correlation in returns  $r_t$  is weak.

**First step: Modeling  $\mu_t$ .** The equation for the mean  $\mu_t$  should be simple. Often we set  $\mu_t = \mu$  constant!

If  $r_t$  are correlated,  $\mu_t$  can be modeled by an ARMA(p,q) model with some explanatory variables  $\xi_{i,t}$ :

$$r_t = \mu_t + r_t^*, \quad \mu_t = \phi_0 + \sum_{i=1}^p \beta_i r_{t-i} - \sum_{i=1}^q \theta_i r_{t-i}^* + \sum_{i=1}^p \phi_i \xi_{i,t-i},$$

where

- $\xi_{i,t}$  are some explanatory variables,
- $p$  and  $q$  is the order of an ARMA model and
- $r_i^*$  are uncorrelated variables (white noise) with 0 mean.

The order  $p, q$  might depend on frequency of the data:

- daily returns  $r_t$  might show some minor correlation,
- monthly returns  $r_t$  tend to be uncorrelated. Then we set  $\mu_t = \mu$ .

A dummy variable  $\xi_{i,t}$  might be for Mondays to study weekend effect.

Combining equations we have

$$\sigma_t^2 = \text{Var}(r_t | F_{t-1}) = \text{Var}(r_t^* | F_{t-1}) = E[(r_t - \mu_t)^2 | F_{t-1}].$$


**Definition:**  $\sigma_t^2$  is called volatility.

We will model evolution of the volatility  $\sigma_t^2$  in time. Such evolution will be governed by

- ARCH -GARCH models.

Besides these models volatility can be modeled by stochastic volatility models where  $\sigma_t^2$  is defined by a stochastic equation.

**Example** Volatility plot. Note: volatility  $\sigma_t^2$  is non-negative, it is not observed directly! We need to estimate/ extract it from data.



### 5.1.2 Model building

Constructing volatility model for a return series consists of three steps:

- Specify a model for a mean  $\mu_t$ . If no correlation in returns  $r_t$ , one can set

$$\mu_t = \mu$$

constant. If returns  $r_t$  correlated, you can fit a model to  $r_t$ . It can be an ARMA model, if needed, to remove dependence.

- Use residuals  $r_t^* = r_t - \mu_t$ , to test for ARCH effects in  $r_t^*$ .

ARCH effect means:

- $r_t^*$  are uncorrelated,
  - the squares  $r_t^{*2}$  correlated
- If ARCH effect is significant, specify volatility model for  $\sigma_t^2$  and perform joint estimation of  $\mu_t$  and  $\sigma_t^2$ .

For most asset return series  $r_t$ , the correlation is weak. So mean equation  $r_t = \mu + r_t^*$  results in removing the sample mean  $\bar{r}$  from the data  $r_t$  if the

sample mean is significantly different from zero:

$$r_t^* = r_t - \bar{r}, \quad \bar{r} = T^{-1} \sum_{j=1}^T r_j.$$

For daily returns, a simple AR(p) model might be needed to be fitted to  $r_t$  to remove correlation in  $r_t$ .

The mean equation may also employ some explanatory variables, such as dummy variables for weekend or January effects.

Poll 2

## 5.2 Testing for ARCH effect

**ARCH effect.** Testing for ARCH effect (or conditional heteroscedasticity) in the series  $r_t = \mu_t + r_t^*$ , means testing for

- no correlation in residuals  $r_t^* = r_t - \mu_t$
- correlation in squares  $r_t^{*2}$

Two tests can be used.

**Ljung-Box test.** We can apply Ljung-Box statistics  $Q(m)$  to the squares  $r_t^{*2}$ . Then the null hypothesis is that the first  $m$  ACF's are zero:

$$\rho_1 = 0, \quad \rho_2 = 0, \quad \dots, \quad \rho_m = 0.$$

**The Lagrange multiplier test** (by Engle (1982)). It is the usual test  $F$  test testing for the null hypothesis

$$H_0: \alpha_1 = \dots = \alpha_m = 0$$

in the linear regression

$$r_t^{*2} = \alpha_0 + \alpha_1 r_{t-1}^{*2} + \dots + \alpha_m r_{t-m}^{*2} + e_t, \quad t = m + 1, \dots, N$$

where  $e_t$  denotes the error term,  $m \geq 1$  is the lag and  $N$  is the sample size.

**Note:** Under  $H_0$ ,

$$r_t^{*2} = \alpha_0 + e_t$$

$\leftarrow \alpha_1 = \alpha_2 = \dots = \alpha_m = 0$

is white noise. Then,  $r_t^{*2}$  are uncorrelated variables.



Denote

$$SSR_0 = \sum_{t=m+1}^N (r_t^{*2} - \bar{\omega})^2, \quad \text{where } \bar{\omega} = (1/N) \sum_{t=1}^N r_t^{*2},$$
$$SSR_1 = \sum_{t=m+1}^N \hat{e}_t^2, \quad \hat{e}_t = r_t^{*2} - \hat{\alpha}_0 - \hat{\alpha}_1 r_{t-1}^{*2} - \dots - \hat{\alpha}_m r_{t-m}^{*2}$$

where  $\hat{e}_t$  is the least squares residual of the previous linear regression.

Then, under  $H_0$ , the statistic

$$F = \frac{(SSR_0 - SSR_1)/m}{SSR_1/(N - 2m - 1)} \sim \chi_m^2$$

has asymptotical chi-squared distribution with with  $m$  degrees of freedom.

*p-value  
will be  
given*

**Example.** Consider the monthly log stock returns of Intel Corporation from 1873 to 2003. The series has no significant correlation so it can be used directly to test for an ARCH effect.

—  $Q(12) = 18.57$  with  $p$ -value 0.10 confirms no serial correlation in  $r_t^{*2}$ . So, no ARCH effect.

— On the other hand, the Lagrange multiplier test shows strong ARCH effect with test statistic  $F \sim 43.5$  and  $p$ -value close to zero.

### ***S-Plus Demonstration***

Denote the return series by `intc`. Note that the command `archTest` applies directly to the  $a_t$  series, not to  $a_t^2$ .

```
> autocorTest(intc, lag=12)
Test for Autocorrelation: Ljung-Box
Null Hypothesis: no autocorrelation
```

```
Test Statistics:
Test Stat 18.5664 p.value 0.0995
```

```
Dist. under Null: chi-square with 12 degrees of freedom
Total Observ.: 372
```

```
> archTest(intc, lag=12)
Test for ARCH Effects: LM Test
Null Hypothesis: no ARCH effects
```

```
Test Statistics:
Test Stat 43.5041 p.value 0.0000
```

```
Dist. under Null: chi-square with 12 degrees of freedom
```

*roll 3  
ARCH  
effect*

**Poll 3** Test for ARCH effect in returns  $r_t$ . Sample size  $N=400$

**Correlogram of  $r_t$**

Date: 29/11/20 Time: 10:53  
 Sample: 1 400  
 Included observations: 400

Autocorrelation	Partial Correlation	AC	PAC	Q-Stat	Prob
		1 -0.023	-0.023	0.2196	0.639
		2 -0.032	-0.033	0.6458	0.724
		3 -0.036	-0.037	1.1600	0.763
		4 -0.057	-0.060	2.4792	0.648
		5 0.050	0.045	3.4877	0.625
		6 -0.076	-0.080	5.8704	0.438
		7 0.067	0.063	7.7087	0.359
		8 -0.009	-0.012	7.7392	0.459
		9 0.012	0.016	7.7940	0.555
		10 -0.041	-0.049	8.4897	0.581

**Correlogram of  $r_t^2$**

Date: 29/11/20 Time: 10:54  
 Sample: 1 400  
 Included observations: 400

Autocorrelation	Partial Correlation	AC	PAC	Q-Stat	Prob
		1 -0.007	-0.007	0.0189	0.891
		2 -0.050	-0.050	1.0282	0.598
		3 -0.026	-0.027	1.3033	0.728
		4 0.025	0.023	1.5643	0.815
		5 -0.023	-0.025	1.7743	0.879
		6 0.060	0.062	3.2521	0.777
		7 0.015	0.015	3.3485	0.851
		8 0.021	0.026	3.5356	0.896
		9 0.026	0.032	3.8035	0.924
		10 -0.053	-0.054	4.9636	0.894

$z_t^*$  is white noise!

### 5.3 The ARCH model

ARCH model was suggested by Engle (1982). It provides a framework for volatility modeling. The idea of such modeling is based on two facts:

- a) the shock  $r_t^*$  of asset return is serially uncorrelated, but dependent,
- b) the dependence of  $r_t^*$  can be described by a simple quadratic function for  $r_t^{*2}$ .

ARCH effect

ARCH(m) model: it assumes that

$$r_t^* = \sigma_t \varepsilon_t, \quad \sigma_t^2 = \alpha_0 + \alpha_1 r_{t-1}^{*2} + \dots + \alpha_m r_{t-m}^{*2}$$

ARCH equation for volatility  $\sigma_t^2$ .

where  $\varepsilon_t$  is a sequence of iid variables with  $E\varepsilon_t = 0$ ,  $E\varepsilon_t^2 = 1$ ,

$$\alpha_0 > 0, \quad \alpha_1 \geq 0, \dots, \alpha_m \geq 0.$$

For existence of a stationary solution, the parameter must satisfy additional conditions. Here,

$$\mu_t = E[r_t^* | F_{t-1}] = 0, \quad \text{var}(r_t^{*2} | F_{t-1}) = \sigma_t^2.$$

In practice,  $\varepsilon_t$  are often assumed to follow the standard-normal  $N(0, 1)$  or Student  $t$ -distribution. In case of this model, the past large shocks  $r_{t-1}^{*2}, \dots, r_{t-m}^{*2}$  implies large conditional variance  $\sigma_t^2$  for the innovation  $r_t^*$ . In ARCH models, large shocks tend to be followed by another large shock. "Tend" means that the probability to obtain large variate is greater than that of smaller variate. So ARCH model has the feature of clustering observed in asset returns.

**Example.** The ARCH effect is rather common in financial time series. Figure 3.2 shows the time plot of

- the percentage change in Deutsche mark/US dollar exchange rate measured in 10 min intervals from June 5, 1989 to June 9, 1989, 2488 observations.
- the squared series of the percentage changes.

We observe, that big changes occur rarely, and there are certain stable periods. Figure 3.3 a) shows that there is no serial correlation in series of percentage changes. Figure 3.3 b) of PACF shows that correlation is present in the squared series of changes. PACF has big spikes, suggesting that series of percentage changes is not independent.

$z_t^2$

↓ implies ARCH effect






Figure 3.2. (a) Time plot of 10-minute returns of the exchange rate between Deutsche mark and U.S. dollar and (b) the squared returns.




Figure 3.3. (a) Sample autocorrelation function of the return series of mark/dollar exchange rate and (b) sample partial autocorrelation function of the squared returns.



## 5.4 Properties of ARCH(1) model

### ARCH(1) model.

To understand ARCH models, first we study ARCH(1) model

$$r_t^* = \sigma_t \varepsilon_t, \quad \sigma_t^2 = \alpha_0 + \alpha_1 r_{t-1}^{*2},$$

with parameters  $\alpha_0 > 0, \alpha_1 \geq 0$ .

Unconditional mean:

$$E[r_t^*] = 0$$

Variance:

$$Var(r_t^*) = \alpha_0 + \alpha_1 Var(r_t^*)$$

and

$$Var(r_t^*) = \frac{\alpha_0}{1 - \alpha_1}$$

Since the variance must be positive, we require

$$0 \leq \alpha_1 < 1.$$

• the kurtosis of  $r_t^*$  is

$$K(r_t^*) = \frac{E(r_t^{*4})}{[Var(r_t^*)]^2} = \frac{3\alpha_0^2(1 + \alpha_1)}{(1 - \alpha_1)(1 - 3\alpha_1^2)} \cdot \frac{(1 - \alpha_1)^2}{\alpha_0^2} = 3 \frac{1 - \alpha_1^2}{1 - 3\alpha_1^2} > 3.$$

It is positive and greater than 3.

$\alpha_1$  must satisfy condition

$$1 - 3\alpha_1^2 > 0,$$

that is  $0 \leq \alpha_1 < 1/3$ ,

### Weaknesses of ARCH models:

1. ARCH models assumes that positive and negative shocks have the same effect on volatility, what is not observed in practise,
2. The ARCH models impose restrictive assumptions on parameters.
3. ARCH models do not explain source of variation.
4. ARCH models are over-predicting the volatility

### 5.4.1 Building ARCH model

Specifying the order of ARCH model is rather easy.

**Order determination.** One can use PACF of  $r_t^{*2}$ . The model we have is

$$\sigma_t^2 = \alpha_0 + \alpha_1 r_{t-1}^{*2} + \dots + \alpha_p r_{t-p}^{*2}.$$

For a given sample,

$r_t^{*2}$  is an unbiased estimator for  $\sigma_t^2$ .

Therefore we might expect  $r_t^{*2}$  to be linearly related to  $r_{t-1}^{*2}, \dots, r_{t-p}^{*2}$  in a manner similar to a linear regression. In general,  $r_t^{*2}$  is not an efficient estimator for  $\sigma_t^2$ , but an approximation, when we identifying order  $p$ .

Alternatively, define

$$\eta_t = r_t^{*2} - \sigma_t^2.$$

*uncorrelated*

It can be shown that  $\eta_t$  is uncorrelated series (white noise) and has zero mean. The ARCH model we can write as an AR(p) model

$$\begin{aligned} r_t^{*2} &= \sigma_t^2 + \eta_t, \\ r_t^{*2} &= \alpha_0 + \alpha_1 r_{t-1}^{*2} + \dots + \alpha_p r_{t-p}^{*2} + \eta_t, \end{aligned}$$

where  $\eta_t$  is a white noise but not an iid.

The PACF function might be useful determining the order of the model. *ARCH(p)*

**Parameter estimation.** Parameters of ARCH model can be estimated using (conditional) Maximum Likelihood method both in case when the noise  $\varepsilon_t$  has normal or Student-t distribution.

**Forecasting.** Forecast of an ARCH model can be obtained recursively as in case of AR models.

Let be given ARCH(p) model. Then the 1-step ahead forecast of volatility is

$$\begin{aligned} \sigma_t^2(1) &= E[\sigma_{t+1}^2 | F_t] = E[\alpha_0 + \alpha_1 r_t^{*2} + \dots + \alpha_p r_{t+1-p}^{*2} | F_t] \\ &= \alpha_0 + \alpha_1 r_t^{*2} + \dots + \alpha_p r_{t+1-p}^{*2}. \end{aligned}$$

The 2-step ahead forecast is

$$\sigma_t^2(2) = \alpha_0 + \alpha_1 \sigma_t^2(1) + \alpha_2 r_t^{*2} + \dots + \alpha_p r_{t+2-p}^{*2}.$$

*Forecasting of volatility*

## 5.5 Examples of ARCH modeling

**Example 3.1.** We shall build an ARCH model for the monthly log returns of Intel stock. The sample ACF and PACF in Figure 3.1 show the presence of conditional heteroscedasticity. It is confirmed by test for ARCH effects.

We proceed to identify the order  $m$  of ARCH model. The PACF in Figure 3.1(d) indicates that an ARCH(3) model might be appropriate. No correlation in  $r_t$ . So we specify the model:

$$r_t = \mu + r_t^*, \quad r_t^* = \sigma_t \varepsilon_t, \quad \sigma_t^2 = \alpha_0 + \alpha_1 r_{t-1}^{*2} + \alpha_2 r_{t-2}^{*2} + \alpha_3 r_{t-3}^{*2}$$

for monthly returns of Intel stock. Assuming that  $\varepsilon_t$  are iid standard normal, we obtain the fitted model:

$$r_t = 0.0171 + r_t^*, \quad \sigma_t^2 = 0.0120 + 0.178r_{t-1}^{*2} + 0.0772r_{t-2}^{*2} + 0.0572r_{t-3}^{*2},$$

where standard errors for parameters are 0.0066, 0.0011, 0.0803, 0.0506 and 0.0769, see output below. The estimates for  $\alpha_2$  and  $\alpha_3$  are statistically non-significant at 5-percent level. Therefore the model can be simplified to

$$r_t = 0.0174 + r_t^*, \quad \sigma_t^2 = 0.0126 + 0.352r_{t-1}^{*2},$$

where the standard errors are 0.0062, 0.0012 and 0.0885, respectively. All estimates are highly significant.

To check goodness of fit we have to investigate residuals


$$\tilde{\varepsilon}_t = \frac{r_t^*}{\sigma_t}.$$

Figure 3.4 shows the residuals and sample ACF of some functions of  $\tilde{\varepsilon}_t$ .


To check that  $\tilde{\varepsilon}_t$  is an iid sequence Ljung-Box statistics  $Q(10)$  gives  $p$ -values 0.18, and for  $\tilde{\varepsilon}_t^2$ , statistics  $Q(10) = 11.38$  has  $p$ -value 0.33, see output. Consequently, ARCH(1) model is an adequate model for describing the data at 5-percent significance level.

Equation above shows that the Intel stock has 1.74 -percent monthly return which is very high. Secondly,  $\hat{\alpha}_1^2 = 0.353^2 < 1/3$ , so that the ARCH(1) model is stationary. and finite fourth moment exists. The unconditional standard deviation is

$$\frac{\alpha_0}{1 - \alpha_1} = \sqrt{0.0126(1 - 0.352)} = 0.1394.$$



**Figure 3.1** Time plot of monthly log returns of Intel stock from January 1973 to December 2008.



**Figure 3.2** Sample ACF and PACF of various functions of monthly log stock returns of Intel Corporation from January 1973 to December 2008: (a) ACF of the log returns, (b) ACF of the squared log returns, (c) ACF of the absolute log returns, and (d) PACF of the squared log returns.



### S-Plus Demonstration

Output edited and % marks explanation.

```
> arch3.fit=garch(intc~1,~garch(3,0))
> summary(arch3.fit)
Call:
garch(formula.mean = intc ~ 1, formula.var = ~ garch(3, 0))

Mean Equation: intc ~ 1
Conditional Variance Equation: ~ garch(3, 0)
Conditional Distribution: gaussian
-----
Estimated Coefficients:
-----
      Value Std.Error t value Pr(>|t|)
  (C) 0.01713  0.006626  2.5860 0.005047 % one-sided
  (A) 0.01199  0.001107 10.8325 0.000000 % p-value
ARCH(1) 0.17874  0.080294  2.2260 0.013309
ARCH(2) 0.07720  0.050552  1.5271 0.063800
ARCH(3) 0.05722  0.076928  0.7438 0.228747
-----
> arch1=garch(intc~1,~garch(1,0)) % A simplified model
> summary(arch1)
Call:
garch(formula.mean = intc ~ 1, formula.var = ~ garch(1,0))

Mean Equation: intc ~ 1
Conditional Variance Equation: ~ garch(1, 0)
Conditional Distribution: gaussian
-----
Estimated Coefficients:
-----
      Value Std.Error t value Pr(>|t|)
  C 0.01741  0.006231  2.794 2.737e-03
  A 0.01258  0.001246 10.091 0.000e+00
ARCH(1) 0.35258  0.088515  3.983 4.094e-05
-----
> stdresi=arch1$residuals/arch1$sigma.t % Standardized
> autocorTest(stdresi,lag=10) % residuals

Null Hypothesis: no autocorrelation
Test Statistics:
Test Stat 13.7820 p.value 0.1832

Dist. under Null: chi-square with 10 degrees of freedom
> archTest(stdresi,lag=10) % ARCH test for residuals

Null Hypothesis: no ARCH effects
Test Statistics:
Test Stat 11.3793 p.value 0.3287

Dist. under Null: chi-square with 10 degrees of freedom
> arch1$asympt.sd % Obtain unconditional variance
[1] 0.1393796

> plot(arch1) % Obtain various plots, including the
% fitted volatility series.
```

Eq (1)


< 0.05

Handwritten notes and markings at the bottom right of the page.

**Poll 5** We would like to fit ARCH model  $r_t = m_t + r^*_t$


We model  $m_t$  by AR(p) model. Select p:

$$r_t = f_1 r_{t-1} + f_2 r_{t-2} + \dots + f_p r_{t-p} + r^*_t$$



**Poll 6** We would like to fit ARCH to  $r^*_t$  which are uncorrelated variables

Select p of ARCH(p) model you would fit to data. Use correlogram



ARCH(1) model can be used for prediction of volatility  $\sigma_t^2$  of Intel stock.

**Example 3.2** Figure 3.2a shows the percentage exchange rate between mark and dollar in 10-minute intervals, we discussed before. Series does not have serial correlation. Sample PACF indicates big spikes at lag 1 and 3. Similarly as in Example 3.1 we specify model ARCH(3) for this data. Unconditional Gaussian maximum likelihood estimation gives the following model:

$$r_t = 0.0018 + r_t^*$$

and

$$r_t^* = \sigma_t \varepsilon_t, \quad \sigma_t^2 = 0.0022 + 0.322r_{t-1}^{*2} + 0.074r_{t-2}^{*2} + 0.093r_{t-3}^{*2}.$$

All estimates are significant at 5% level. Model check for residuals  $\tilde{\varepsilon}_t$  indicates that the estimated ARCH(3) model is adequate.






Figure 3.2. (a) Time plot of 10-minute returns of the exchange rate between Deutsche mark and U.S. dollar and (b) the squared returns.





Fitting ARCH(p) — often too many parameters needed

## 5.6 The GARCH model

The drawback of ARCH(p) model is that often in applications, to describe adequately model, it requires many parameters. We shall have later an example, where the model ARCH(9) is needed for modeling stochastic volatility.

Let  $r_t$  be a log returns series, and

$$r_t = \mu_t + r_t^*$$

where

$$\mu_t = E[r_t | F_{t-1}], \quad r_t^* = r_t - \mu_t$$

is the conditional mean.

### GARCH(p,q) model

We say that the innovations  $a_t$  follow GARCH(p,q) model if

$$r_t^* = \sigma_t \varepsilon_t, \quad \sigma_t^2 = \alpha_0 + \sum_{i=1}^p \alpha_i r_{t-i}^{*2} + \sum_{j=1}^q \beta_j \sigma_{t-j}^2 \quad (20)$$

where  $\varepsilon_t$  are i.i.d. random variables with mean 0 and variance 1,

$$\alpha_0 \geq 0, \alpha_1 \geq 0, \dots, \alpha_p \geq 0, \quad \beta_1 \geq 0, \dots, \beta_q \geq 0 \text{ are parameters.}$$

**Stationary solution.** To assure existence of a stationary solution with a finite variance we impose condition

$$\alpha_1 + \dots + \alpha_p + \beta_1 + \dots + \beta_q < 1. \quad (21)$$

In applications often it is assumed that errors  $\varepsilon_t$  have standard normal or standardized Student- $t$  distribution.

When  $q = 0$ , the GARCH(p,0) model reduces to ARCH(p) model.

$$\text{GARCH}(p,0) = \text{ARCH}(p)$$

## ARMA type representation of GARCH model.

We set

$$\eta_t = r_t^{*2} - \sigma_t^2.$$

← white noise  
(can be shown)

Then  $\eta_t$  are uncorrelated but dependent variables. Then, using  $\sigma_t^2 = r_t^{*2} - \eta_t$ , we obtain,

$$\begin{aligned} r_t^{*2} &= \sigma_t^2 + \eta_t = \alpha_0 + \sum_{i=1}^p \alpha_i r_{t-i}^{*2} + \sum_{j=1}^q \beta_j \sigma_{t-j}^2 + \eta_t \\ &= \alpha_0 + \sum_{i=1}^p \alpha_i r_{t-i}^{*2} + \sum_{j=1}^q \beta_j r_{t-j}^{*2} + \eta_t - \sum_{j=1}^q \beta_j \eta_{t-j}. \end{aligned}$$

ARMA model

Equation is an ARMA form for squares  $r_t^{*2}$  where  $\eta_t$  plays role of a white noise. Using formula for unconditional mean of an ARMA model, we have

$$E[r_t^{*2}] = \frac{\alpha_0}{1 - \alpha_1 - \dots - \alpha_p - \beta_1 - \dots - \beta_q}$$

where condition (21) implies that the denominator is positive and therefore variance  $Var(r_t^{*2})$  is finite.

## GARCH(1,1) model

The simple GARCH model is GARCH (1,1) model where condition variance is modeled by

$$\sigma_t^2 = \alpha_0 + \alpha_1 r_{t-1}^{*2} + \beta_1 \sigma_{t-1}^2, \quad 0 \leq \alpha_1, \beta_1 \leq 1, \quad \alpha_1 + \beta_1 < 1.$$

POLLZ

Properties:

- A large  $r_{t-1}^{*2}$  or  $\sigma_{t-1}^2$  leads to a large  $\sigma_t^2$ . This means that large  $r_{t-1}^{*2}$  is followed by another large  $r_t^{*2}$ , so we have clustering property, common in financial time series.

- The excess kurtosis is greater than 3:

under condition  $1 - 2\alpha_1^2 - (\alpha_1 + \beta_1)^2 > 0$ , we have

$$\frac{E[r_t^{*4}]}{(E[r_t^{*2}])^2} = \frac{3[1 - (\alpha_1 + \beta_1)^2]}{1 - (\alpha_1 + \beta_1)^2 - 2\alpha_1^2} > 3.$$

This shows that the tail of GARCH(1,1) distribution is heavier than that of a normal distribution.

- GARCH(1,1) model is a simple parametrical model which can be used to describe volatility evolution.

**Forecasting of GARCH(1,1) model.** Forecasting of GARCH models is similar to that of ARMA models. To obtain 1-step ahead forecast of volatility  $\sigma_t^2$ , note that

$$\sigma_{t+1}^2 = \alpha_0 + \alpha_1 r_t^{*2} + \beta_1 \sigma_t^2,$$

*volatility*

where  $r_t^{*2}, \sigma_t^2$  are known at time  $t$  and parameters  $\alpha_0, \alpha_1, \beta_1$  are estimated from the data.

**1-step ahead forecast** at horizon  $t$  is

$$\sigma_t^2(1) = E[\sigma_{t+1}^2 | F_t] = \alpha_0 + \alpha_1 r_t^{*2} + \beta_1 \sigma_t^2.$$

**2-step ahead forecast.** We rewrite the volatility equation as

$$\sigma_{t+2}^2 = \alpha_0 + (\alpha_1 + \beta_1) r_{t+1}^{*2} + \alpha_1 \sigma_{t+1}^2 (\varepsilon_{t+1}^2 - 1).$$

Since  $E(\varepsilon_{t+1}^2 - 1 | F_t) = 0$  then

$$\sigma_t^2(2) = E[\sigma_{t+2}^2 | F_t] = \alpha_0 + (\alpha_1 + \beta_1) \sigma_t^2(1).$$

**k-step ahead prediction:** for  $k \geq 2$ , is

$$\sigma_t^2(k) = E[\sigma_{t+k}^2 | F_t] = \alpha_0 + (\alpha_1 + \beta_1) \sigma_t^2(k-1).$$

By repeated substitutions in the above equation we obtain the formula

$$\sigma_t^2(k) = \frac{\alpha_0 [1 - (\alpha_1 + \beta_1)^{k-1}]}{1 - \alpha_1 - \beta_1} + (\alpha_1 + \beta_1)^{k-1} \sigma_t^2(1).$$

We see that

$$\sigma_t^2(k) \rightarrow \frac{\alpha_0}{1 - \alpha_1 - \beta_1} = E[\sigma_t^2] = E[r_t^{*2}]$$

*Mean reversion property*

if  $\alpha_1 + \beta_1 < 1$ , and therefore GARCH(1,1) process is mean reverting.

The literature on GARCH models is very rich.

The weakness of GARCH model is the same as in case of ARCH model:

- The model responds equally to positive and negative shocks
- The tails of distribution of GARCH models are not sufficiently heavy comparing to some financial data.

## 5.7 Illustrative example

The modeling procedure used in ARCH case can be used also for GARCH models. However, specifying the order of GARCH models is not an easy task.

In practise, only the lower order models are used:

GARCH(1,1), GARCH(1,2), GARCH(2,1).

Parameters can be estimated using conditional maximum likelihood method assuming that that initial value  $\sigma_1^2$  of volatility is fixed (known). Then, in GARCH(1,1) case, the volatility  $\sigma_t^2$  can be computed recursively.

- In some applications a good starting value of  $\sigma_1^2$  might be the sample variance of  $r_t^{*2}$ . : To compute  $\sigma_t^2$
- The fitted model, its goodness, can be checked by using standardized residuals

$$\tilde{\varepsilon}_t = \frac{r_t^*}{\sigma_t}$$

and its squared model

$$\tilde{\varepsilon}_t^2 = \frac{r_t^{*2}}{\sigma_t^2}$$

If case of good fit, there should be no ARCH effect in residuals  $\tilde{\varepsilon}_t$ , i.e.

- $\tilde{\varepsilon}_t$  should be uncorrelated,
- $\tilde{\varepsilon}_t^2$  should be uncorrelated.

**Example** We shall discuss the monthly excess returns of and S&P index from 1926 for 792 observations. Denote excess return  $r_t$

Definition: Excess return of an asset at time  $t$  is the difference between the assets return and some reference asset:

$$r_t = r_{t,asset} - r_{t,ref.asset}$$

The reference asset is often taken to be a riskless such as a short-term US treasury bill returns.






Figure 3.5. Time series plot of the monthly excess returns of the S&P 500 index.




Figure 3.6. (a) Sample ACF of the monthly excess returns of the S&P 500 index and (b) sample PACF of the squared monthly excess returns.

In Figure 3.6 we see the sample ACF of  $r_t$ , and the sample PACF of  $r_t^2$ :

- the ACF has serial correlation at lags 1 and 3;
- the PACF shows strong linear dependence (correlation) for a number of lags. That means that for  $r_t^2$  ARCH models do not fit well, and we shall fit GARCH(1,1) model.

**Model** The ACF suggests to fit for  $r_t$  MA(3) model. The estimated model:

$$r_t = 0.0062 + r_t^* + 0.0944r_{t-1}^* - 0.1407r_{t-3}^* + \varepsilon_t \quad \hat{\sigma}_{r^*}^2 = 0.0576.$$

where all coefficients are significant at 5 % level, and  $\varepsilon_t$  in case of 'good' fit should be a white noise.

To simplify dependence on  $r_t^*$ , instead of MA(3) model we shall fit to the data an AR(3) model

$$r_t = \phi_1 r_{t-1} + \phi_2 r_{t-2} + \phi_3 r_{t-3} + \phi_0 + r_t^*$$

where  $\phi_1, \phi_2, \phi_3, \phi_0$  are unknown parameters and  $ar_t^*$  is a white noise. The fitted model, assuming normality of  $r_t^*$  is

$$r_t = 0,088r_{t-1} - 0,023r_{t-2} - 0,123r_{t-3} + 0,0066 + r_t^*, \quad \hat{\sigma}_{r^*}^2 = 0,00333.$$

Although the residuals  $r_t^*$  might be a white noise (uncorrelated), they still might be dependent. To model this type of data (so called GARCH effect), we use GARCH(1,1) model for  $\sigma_t$ :

$$r_t^* = \sigma_t \varepsilon_t, \quad \sigma_t^2 = \alpha_0 + \beta_1 \sigma_{t-1}^2 + \alpha_1 r_{t-1}^{*2}.$$

A joint estimation of the AR(3)-GARCH(1,1) model gives

$$r_t = 0.0078 + 0.032r_{t-1} - 0.029r_{t-2} - 0.008r_{t-3} + r_t^*, \quad \text{AR(3)}$$

$$\sigma_t^2 = 0.000084 + 0.121r_{t-1}^{*2} + 0.8523\sigma_{t-1}^2. \quad \text{GARCH(1,1)}$$

Having estimates of parameters we can estimate the unconditional variance of  $r_t^*$ :

$$Var(r_t^*) \sim \frac{\alpha_0}{1 - (\alpha_1 + \beta_1)} = \frac{0.000084}{1 - 0.8523 - 0.1213} = 0.00317$$

which is close to that derived fitting AR(3) model to  $r_t$ .

**t-ratios** suggest that all three parameter in the mean (AR(3) part) are not significant. We refine the model, dropping those parameters. The estimated model then is

$$r_t = 0.0076 + r_t^*, \quad \sigma_t^2 = 0.000086 + 0.1216r_{t-1}^{*2} + 0.8511\sigma_{t-1}^2.$$

GARCH(1,1)

The standard error of the constant in the mean equation is 0.0015, so parameter 0.0076 is significant at 5% level.

The standard errors of the parameters in the volatility equation are 0.000024, 0.0197, and 0.0190, respectively. These parameter are significant. The unconditional variance of  $r_t^*$  is

$$\frac{0.000086}{1 - 0.8511 - 0.1216} = 0.00314.$$

This is a simple GARCH(1,1) model.

**Goodness-of fit** To check the goodness of fit of the estimated model we compute standardized shocks

$$\tilde{\varepsilon}_t = \frac{r_t^*}{\sigma_t}.$$

- Figure 3.7 shows the estimated volatility  $\sigma_t^2$  and the standardized shocks  $\tilde{\varepsilon}_t$ .

- Figure 3.8 shows the sample ACF of the residuals  $\tilde{\varepsilon}_t$  and squares  $\tilde{\varepsilon}_t^2$ . The ACF does not show any correlation in  $\tilde{\varepsilon}_t$  and  $\tilde{\varepsilon}_t^2$ , which indicates a good fit of the model.

The Ljung-Box statistics takes value

-  $Q(12) = 11.99$  with  $p$ -value 0.45 for  $\tilde{\varepsilon}_t$ , and

-  $Q(12) = 11.11$  with  $p$ -value 0.36 for  $\tilde{\varepsilon}_t^2$ .

The model appears to be adequate, describing the dependence structure in the return and volatility series.

**Note:** The fitted model has property

$$\hat{\alpha}_1 + \hat{\beta}_1 = 0.9772$$

is close to 1. This phenomenon is often observed in practice and leads to the model GARCH(1,1) with constrain  $\alpha_1 + \beta_1 = 1$ . The GARCH(1,1) model with this property is called IGARCH model.




Figure 3.7. (a) Time series plot of estimated volatility ( $\sigma_t$ ) for the monthly excess returns of the S&P 500 index and (b) the standardized shocks of the monthly excess returns of the S&P 500 index. Both plots are based on the GARCH(1,1) model in Eq. (3.18).




Figure 3.8. Model checking of the GARCH(1,1) model in Eq. (3.18) for monthly excess returns of the S&P 500 index: (a) sample ACF of standardized residuals and (b) sample ACF of the squared standardized residuals.



**Forecasting:** To forecast the volatility of monthly excess returns of the S&P index, we use the volatility equation

$$r_t = 0.0076 + r_t^*, \quad \sigma_t^2 = 0.000086 + 0.1216r_{t-1}^{*2} + 0.8511\sigma_{t-1}^2.$$

GARCH(1,1)

Then the 1-set ahead forecast is

$$\sigma_t^2(1) = 0.00008 + 0.121r_t^{*2} + 0.8511\sigma_t^2$$

where  $r_t^{*2}$  is the residual, in our case  $r_t^* = r_t - 0.0076$ , and  $\sigma_t^2$  can be computed using GARCH(1,1) equation for volatility. Computing it, we can fix  $\sigma_0^2$  at zero or at unconditional variance of  $r_t^*$ . For multi-step forecast, we use recursive formula.

Table 3.2 shows some mean and volatility forecasts for the returns of the S&P index with forecast horizon  $t = 792$ .

**Table 3.1. Volatility Forecasts for the Monthly Excess Returns of the S&P 500 Index<sup>a</sup>**

Horizon	1	2	3	4	5	$\infty$
Return	0.0076	0.0076	0.0076	0.0076	0.0076	0.0076
Volatility	0.0536	0.0537	0.0537	0.0538	0.0538	0.0560

<sup>a</sup>The forecast origin is  $h = 792$ , which corresponds to December 1991. Here volatility denotes conditional standard deviation.

excess return of the S&P 500 index with forecast origin  $h = 792$  based on the GARCH(1,1) model in Eq. (3.18).

**Some S-Plus Commands Used in Example 3.3**

```
> fit=garch(sp-ar(3),~garch(1,1))
> summary(fit)
> fit=garch(sp-1,~garch(1,1))
> summary(fit)
> names(fit)
[1] "residuals" "sigma.t" "df.residual" "coef" "model"
[6] "cond.dist" "likelihood" "opt.index" "cov"
[10] "prediction" "call" "asympt.sd" "series"
> % Next, compute the standardized residuals
> stdresi=fit$residuals/fit$sigma.t
> autocorTest(stdresi,lag=24)
> autocorTest(stdresi^2,lag=24)
> predict(fit,5) % Compute predictions
```

Note that in the prior commands the volatility series  $\sigma_t$  is stored in `fit$sigma.t` and the residual series of the returns in `fit$residuals`.

**S-Plus Commands Used**

```
> fit1 = garch(sp-1,~garch(1,1),cond.dist='t',cond.par=5,
+ cond.est=F)
> names(fit1)
```