

MTH6107 Chaos & Fractals

Solutions 2

Exercise 1. Order the integers from 1 to 50 inclusive using Sharkovskii's ordering.

1 \triangleleft 2 \triangleleft 4 \triangleleft 8 \triangleleft 16 \triangleleft 32 \triangleleft 48 \triangleleft 40 \triangleleft 24 \triangleleft 44 \triangleleft 36 \triangleleft 28 \triangleleft 20 \triangleleft 12 \triangleleft 50 \triangleleft 46 \triangleleft 42 \triangleleft 38 \triangleleft 34 \triangleleft 30 \triangleleft 26 \triangleleft 22 \triangleleft 18 \triangleleft 14 \triangleleft 10 \triangleleft 6 \triangleleft 49 \triangleleft 47 \triangleleft 45 \triangleleft 43 \triangleleft 41 \triangleleft 39 \triangleleft 37 \triangleleft 35 \triangleleft 33 \triangleleft 31 \triangleleft 29 \triangleleft 27 \triangleleft 25 \triangleleft 23 \triangleleft 21 \triangleleft 19 \triangleleft 17 \triangleleft 15 \triangleleft 13 \triangleleft 11 \triangleleft 9 \triangleleft 7 \triangleleft 5 \triangleleft 3

Exercise 2. For the map $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = (x - 1)(1 - 3x^2/2)$, determine the orbit of the point 0.

It is the period-3 orbit $\{0, -1, 1\}$.

Exercise 3. Use Sharkovskii's Theorem to show that the map $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = (x - 1)(1 - 3x^2/2)$ has a point of minimal period n for every $n \in \mathbb{N}$.

The map f is certainly continuous, so the existence (see Exercise 2) of an orbit of minimal (i.e. prime) period 3 implies, by Sharkovskii's Theorem, the existence of points of minimal period n for all $n \in \mathbb{N}$.

Exercise 4. Give an example of a continuous map $f : \mathbb{R} \rightarrow \mathbb{R}$ which has one fixed point, and no other periodic points.

We might choose $f(x) = 2x$ (or indeed $f(x) = cx$ for any $c \neq 1$). Alternatively, we might choose f to be any order reversing diffeomorphism.

Exercise 5. Give an example of a continuous map $f : \mathbb{R} \rightarrow \mathbb{R}$ which has three fixed points, and no other periodic points.

One such example is $f(x) = x^3$. Note that the three fixed points are at $-1, 0$, and 1 . To see that there are no other periodic points, note that if $x \in (0, 1)$ or $x \in (-1, 0)$ then $f^n(x)$ converges to 0 , if $x \in (1, \infty)$ then $f^n(x)$ converges to ∞ , while if $x \in (-\infty, -1)$ then $f^n(x)$ converges to $-\infty$; therefore if $x \in \mathbb{R} \setminus \{-1, 0, 1\}$ then x is not periodic.

Exercise 6. Give an example of a continuous map $f : \mathbb{R} \rightarrow \mathbb{R}$ which has one fixed point, one orbit of prime period two, and no other periodic points.

One such example is $f(x) = -x^3$. Note that the unique fixed point is at 0 (i.e. the unique real solution of $x^3 + x = 0$). The points -1 and 1 are of prime period 2 (together with the point 0 they are the only real solutions of $0 = x^9 - x = x(x + 1)(x - 1)(x^2 + 1)(x^4 + 1)$). To see that there are no other periodic points, note that if $x \in (0, 1)$ or $x \in (-1, 0)$ then $f^n(x)$ converges to 0 , while if $x \in (1, \infty)$ or $x \in (-\infty, -1)$ then $|f^n(x)|$ converges to ∞ ; therefore if $x \in \mathbb{R} \setminus \{-1, 0, 1\}$ then x is not periodic.

Exercise 7. Give an example of a map $f : \mathbb{R} \rightarrow \mathbb{R}$ which has one orbit of minimal period three, and no other periodic points.

Note that the question does not ask for f to be a *continuous* map (in fact a continuous map with the required properties does not exist, by Sharkovskii's Theorem). However we can easily exhibit a discontinuous map with these properties: for example let f be the map defined by setting $f(-1) = 0$, $f(0) = 1$, $f(1) = -1$, and $f(x) = 0$ for all $x \in \mathbb{R} \setminus \{-1, 0, 1\}$. Then $\{-1, 0, 1\}$ is an orbit of minimal period 3, and all points in $\mathbb{R} \setminus \{-1, 0, 1\}$ are preperiodic but not periodic.

Exercise 8. For the following values of μ , describe the behaviour of the orbit of the point x_0 under the logistic map $f_\mu(x) = \mu x(1 - x)$.

- (a) $\mu = 4/5 = 0.8$, $x_0 = 3/5 = 0.6$,
- (b) $\mu = 7/5 = 1.4$, $x_0 = 1/2 = 0.5$,
- (c) $\mu = 33/10 = 3.3$, $x_0 = 13/20 = 0.65$,
- (d) $\mu = 4$, $x_0 = 13/20 = 0.65$,
- (e) $\mu = 4$, $x_0 = 33/50 = 0.66$.

(a) Letting $x_i = f_{0.8}^i(x_0)$ then $x_0 = 0.6$, $x_1 = 0.192$, $x_2 = 0.124109$, $x_3 = 0.869646$, $x_4 = 0.0635214$, etc., and $x_i \rightarrow 0$ as $i \rightarrow \infty$.

(b) Letting $x_i = f_{1.4}^i(x_0)$ then $x_0 = 0.5$, $x_1 = 0.35$, $x_2 = 0.3185$, $x_3 = 0.303881$, $x_4 = 0.296152$, $x_5 = 0.291824$, $x_6 = 0.289328$, etc., and $x_i \rightarrow (\mu - 1)/\mu = 2/7 = 0.285714 \dots$ as $i \rightarrow \infty$.

(c) Letting $x_i = f_{3.3}^i(x_0)$ then $x_0 = 0.65$, $x_1 = 0.75075$, $x_2 = 0.617511$, $x_3 = 0.779431$, $x_4 = 0.56733$, $x_5 = 0.81004$, $x_6 = 0.507788$, $x_7 = 0.8248$, $x_8 = 0.476867$, etc., and the orbit is in the basin of attraction of the attracting 2-cycle $\{0.479427, 0.823603\}$.

(d) Letting $x_i = f_4^i(x_0)$ then $x_0 = 0.65$, $x_1 = 0.91$, $x_2 = 0.3276$, $x_3 = 0.881113$, $x_4 = 0.419012$, $x_5 = 0.973764$, $x_6 = 0.102192$, and x_i does not seem to settle to any discernible pattern.

(e) Letting $x_i = f_4^i(x_0)$ then $x_0 = 0.66$, $x_1 = 0.8976$, $x_2 = 0.367657$, $x_3 = 0.929941$, $x_4 = 0.260602$, $x_5 = 0.770754$, $x_6 = 0.706768$, and x_i does not seem to settle to any discernible pattern. Moreover note that this orbit of $x_0 = 0.66$ is already looking rather different to the orbit of 0.65 in (d) above; this is a hallmark of *chaos*, that two orbits which start close to each other end up looking nothing like each other.

Exercise 9. Let $f(x) = 1 - (13/10)x^2 = 1 - 1.3x^2$.

Use a computer to determine numerically the first 50 points in the f -orbit of the point 0, and the first 50 points in the f -orbit of the point $1/3$.

What is the period of the attracting orbit of f ?

The points $f^n(0)$ for $0 \leq n \leq 50$, listed in order, and to six-digit precision, are: 0, 1, -0.3, 0.883, -0.0135957, 0.99976, -0.299375, 0.883487, -0.0147135, 0.999719, -0.299268, 0.88357, -0.0149047, 0.999711, -0.299249, 0.883585, -0.0149389, 0.99971, -0.299246, 0.883588, -0.014945, 0.99971, -0.299245, 0.883588, -0.0149461, 0.99971, -0.299245, 0.883588, -0.0149463, 0.99971, -0.299245, 0.883588, -0.0149464, 0.99971, -0.299245, 0.883588, -0.0149464, 0.99971, -0.299245, 0.883588, -0.0149464, 0.99971,

-0.299245, 0.883588, -0.0149464, 0.99971, -0.299245, 0.883588, -0.0149464, 0.99971, -0.299245.

The points $f^n(1/3)$ for $0 \leq n \leq 50$, listed in order, and to six-digit precision, are: 0.333333, 0.855556, 0.0484321, 0.996951, -0.292084, 0.889093, -0.0276328, 0.999007, -0.29742, 0.885003, -0.0182004, 0.999569, -0.298881, 0.883872, -0.0155975, 0.999684, -0.299178, 0.88364, -0.0150665, 0.999705, -0.299233, 0.883598, -0.0149681, 0.999709, -0.299243, 0.88359, -0.0149503, 0.999709, -0.299245, 0.883588, -0.0149471, 0.99971, -0.299245, 0.883588, -0.0149465, 0.99971, -0.299245, 0.883588, -0.0149464, 0.99971, -0.299245, 0.883588, -0.0149464, 0.99971, -0.299245, 0.883588, -0.0149464, 0.99971, -0.299245, 0.883588, -0.0149464.

We observe that there is an attracting period-4 orbit. (Note that the orbits of the points 0 and $1/3$ get very close to, but do not ever reach, the attracting period-4 orbit: this can be seen e.g. by using higher precision calculations).

Exercise 10. Let $f(x) = 1 - \bar{\lambda}x^2$ where the value

$$\bar{\lambda} = \frac{1}{3} \left(2 + \left(25/2 - 3\sqrt{69}/2 \right)^{1/3} + \left(25/2 + 3\sqrt{69}/2 \right)^{1/3} \right) \approx 1.75487766624669276$$

is the only real root of the polynomial $1 - \lambda + 2\lambda^2 - \lambda^3$. (The value $\bar{\lambda}$ is chosen so that 0 is a period-3 point).

Use a computer to determine numerically the first 100 points in the f -orbit $\{f^n(1/3)\}_{n=0}^{\infty}$ of the point $1/3$.

What is the smallest value $n \in \mathbb{N}$ such that $|f^n(1/3)| < 1/100$?

What is the smallest value $n \in \mathbb{N}$ such that $|f^n(1/3)| < 1/1000$?

What is the smallest value $n \in \mathbb{N}$ such that $|f^n(1/3)| < 10^{-6}$?

The points $f^n(1/3)$ for $1 \leq n \leq 100$, listed in order, and to 30-digit precision, are: 0.805013592639256359994499011516, -0.13724300399842469417029663821, 0.96694575226767184768572001197, -0.64078269402626004164645869879, 0.27944291154913380895156940089, 0.86296451470939636812027419337, -0.3068710047569593686982052890, 0.8347434973479832615779149072, -0.2227929779137414332615638862, 0.9128936444266934943780011606, -0.462470834722877354274795755, 0.624668040593139431299310199, 0.315228903426547703378679718, 0.825619102184820392690376406, -0.19620692439734415711846210, 0.93244219864825837155853729, -0.52577567356143373447667293, 0.51488151456471208844976748, 0.5347767116008343248170330, 0.4981293253769257634425915, 0.5645572474949496531308910, 0.440676816409665405875317, 0.659209677538190472251826, 0.237404925862833321711402, 0.901093166428749540446619, -0.42490607879351239322009, 0.68316530324758072393065, 0.18097256558688413584035, 0.94252587959735727599594, -0.55895440835686168480563, 0.4517235169974377326302, 0.6419099543643334229536, 0.2769054838324370250360, 0.8654418646983001916738, -0.314385158446140295665, 0.826551352348052206649, -0.198909850463220240671, 0.930568039437850485450, -0.51964838166587839212, 0.52612259112985583610, 0.51424112111767277215, 0.53593325213070198278, 0.4959562261974700898,

