

# Chapter 1

## Powers and Logarithms

### 1.1 Powers or Indices

If  $a \in \mathbb{R}$ , i.e.  $a$  is a real number, and  $n$  is a positive integer  $1, 2, 3, \dots$ , i.e.  $n \in \mathbb{N}$ , then  $a^n$  is a short-hand notation for **multiplying  $a$  by itself  $n$  times**:

$$a^n = \overbrace{a \times \dots \times a}^{n \text{ times}}. \quad (1.1)$$

The number  $a$  is called the **base** and the number  $n$  is referred to as the **power, index** or **exponent**. The operation of raising a number (base) to a power is called **exponentiation**.

For any **base**  $a$  (including zero) and any values of  $m$  and  $n$  the following hold:

Rules for powers			
<b>Rule 1:</b>	$a^1 = a,$		(1.2)
<b>Rule 2:</b>	$a^0 = 1,$		(1.3)
<b>Rule 3:</b>	$a^m \times a^n = a^{m+n},$		(1.4)
<b>Rule 4:</b>	$(a^m)^n = a^{mn},$		(1.5)
<b>Rule 5:</b>	$a^{-1} = \frac{1}{a}$	$(a \neq 0)$	(1.6)
<b>Rule 6:</b>	$a^{-n} = \frac{1}{a^n}$	$(a \neq 0)$	(1.7)
<b>Rule 7:</b>	$a^{1/n} = \sqrt[n]{a}$	$(n \in \mathbb{N}, a > 0)$	(1.8)
<b>Rule 8:</b>	$a^{m/n} = \sqrt[n]{a^m} = (\sqrt[n]{a})^m$	$(n \in \mathbb{N}, a > 0)$	(1.9)
<b>Rule 9:</b>	$(ab)^n = a^n b^n$		(1.10)

#### Remarks:

- 1) Rules 1, 2, 5 and 7 are in fact definitions so that the rules of powers work nicely.
- 2) Observe that  $a$  is typically given in these expressions, while  $m$  and  $n$  take different values.

3) Rules 7 and 8 may fail when  $a < 0$ .  $\sqrt{-1}$  is not defined in  $\mathbb{R}$ , while  $\sqrt[3]{-1} = -1$ .

Let's take a closer look at Rules 3 and 4 when  $m$  and  $n$  are positive integer  $1, 2, 3, \dots$  :

$$a^m \times a^n = \underbrace{a \times \dots \times a}_{m \text{ times}} \times \underbrace{a \times \dots \times a}_{n \text{ times}} = \underbrace{a \times \dots \times a}_{m+n \text{ times}}$$

and

$$(a^m)^n = \underbrace{\underbrace{a \times \dots \times a}_{m \text{ times}} \times \dots \times \underbrace{a \times \dots \times a}_{m \text{ times}}}_{n \text{ times}} = \underbrace{a \times \dots \times a}_{m \times n \text{ times}}.$$

While we have demonstrated how Rules 3 and 4 work when  $m$  and  $n$  take positive integer values we can in fact substitute any fractional or real values into both rules. The same is true for Rule 6, while for Rules 7 and 8 we will only use integer values for  $m$  and  $n$ .

While these worked examples may appear trivial they are good practice the rules of powers.

**Worked Example 1.1.** Using the rules for power show that

1.  $a \times \frac{1}{a} = 1$

2.  $a^n \times \frac{1}{a^n} = 1$

3.  $(\sqrt[n]{a})^n = a$

4.  $(\sqrt[n]{a^m})^n = a^m$

**Solution :**

1.  $\frac{1}{a} = a^{-1}$   $a = a^1$  so  $a \times \frac{1}{a} = a^1 \times a^{-1} = a^{(1+(-1))} = a^0 = 1$   
Rule 5 Rule 1 Rule 3

2.  $\frac{1}{a^n} = a^{-n}$  so  $a^n \times \frac{1}{a^n} = a^n \times a^{-n} = a^{(n+(-n))} = a^0 = 1$   
Rule 6 Rule 3

3.  $\sqrt[n]{a} = a^{1/n}$  so  $(\sqrt[n]{a})^n = (a^{1/n})^n = a^{n \cdot \frac{1}{n}} = a^1 = a$   
Rule 7 Rule 4 Rule 1

4.  $\sqrt[n]{a^m} = a^{m/n}$  so  $(\sqrt[n]{a^m})^n = (a^{m/n})^n = a^{\frac{m}{n} \cdot n} = a^m$   
Rule 8 Rule 4

**Common mistakes**

You need to remember also that, in general,

$a^m + a^n \neq a^{m+n}$	e.g. $3^1 + 3^1 = 3 + 3 = 6 \neq 3^{1+1} = 3^2 = 9$
$(a + b)^n \neq a^n + b^n$	e.g. $(3 + 4)^2 = 7^2 = 49 \neq 3^2 + 4^2 = 25$
$a^{(m^n)} \neq (a^m)^n$	e.g. $2^{(3^2)} = 2^9 = 512 \neq (2^3)^2 = 8^2 = 64.$

**Worked Example 1.2.** Without using a calculator find

i)  $2^6$ ,    ii)  $8^{2/3}$ ,    iii)  $\sqrt[3]{1,000,000}$ ,    iv)  $(0.2)^{-2}$

**Solution :**

$$i) \quad 2^6 = \underbrace{2 \cdot 2 \cdot 2}_8 \cdot \underbrace{2 \cdot 2 \cdot 2}_8 = 8 \cdot 8 = 64$$

$$ii) \quad 8^{2/3} = \sqrt[3]{8^2} = \left(\sqrt[3]{8}\right)^2 = \left(\sqrt[3]{2^3}\right)^2 = 2^2 = 4$$

$$iii) \quad \sqrt[3]{\underbrace{1\,000\,000}_{6 \text{ zeros}}} = \sqrt[3]{10^6} = 10^{6/3} = 10^2 = 100$$

$$iv) \quad (0.2)^{-2} = \left(\frac{1}{5}\right)^{-2} = \left(5^{-1}\right)^{-2} = 5^2 = 25$$

**Worked Example 1.3.** Simplify

$$\frac{\sqrt[7]{a^{22}}}{(a^2 a^3 \sqrt{a})^4}$$

State the rules of powers that you are using at each step.

**Solution :** We simplify bit by bit:

**Numerator:** Rule 8 :  $\sqrt[7]{a^{22}} = a^{22/7}$

**Denominator:**

$$a^2 a^3 \cdot \sqrt{a} = a^{2+3} \cdot \sqrt{a} = a^5 \cdot a^{1/2} = a^{5 + \frac{1}{2}} = a^{11/2}$$

so  $(a^{11/2})^4 = a^{22}$  (Rule 4)

**Overall fraction:**

$$\frac{a^{22/7}}{a^{22}} = a^{22/7 - 22} = a^{-132/7} = \frac{1}{a^{132/7}}$$

(Rule 6, Rule 3, Rule 5)

**Worked Example 1.4.** Calculate the value of

$$\frac{\sqrt[4]{10^{17}}}{(10^2)^2} \times 10^{3/4}$$

Show your work and state the rules you use.

**Solution :** Again we simplify bit by bit:

**Numerator:** Rule 8  $\sqrt[4]{10^{17}} = 10^{17/4}$

**Denominator:**

Rule 4  $(10^2)^2 = 10^4$

**Overall expression:**

$$\frac{10^{17/4}}{10^4} \times 10^{3/4} = \frac{10^{17/4} \cdot 10^{3/4}}{10^4} = \frac{10^{20/4}}{10^4} = \frac{10^5}{10^4} = 10$$

(Rule 3, Rule 6, Rule 3)

Observe that expressions may contain **different bases**, e.g.  $a^m b^n$ . In these cases you **can not apply the rules of powers** until you have transformed all terms to have the same common base using Rule 4. Typically we change to a smaller base.

**Example:**  $9^3 = (3^2)^3 = 3^6$  by Rule 4; and  $1000^4 = (10^3)^4 = 10^{12}$  by Rule 4.

**Worked Example 1.5.** Change  $4^7$  and  $32^2$  to the base 2 and hence find  $\frac{4^7}{32^2}$ .

$$\text{Solution: } 4^7 = (2^2)^7 = 2^{14}$$

$$32^2 = (2^5)^2 = 2^{10}$$

$$\text{Overall: } \frac{4^7}{32^2} = \frac{2^{14}}{2^{10}} = 2^4$$

Rule 3 & 6

**Worked Example 1.6.** Show that

$$(100^6)^{12} \times [10^{-8} \times (0.1)^6]^{10} = 10000$$

**Hint:** Write everything in powers of 10.

**Solution:** Note that a calculator can not help you here (you can try it to see what happens).

$$\text{First term: } (100^6)^{12} = ((10^2)^6)^{12} = (10^{2 \cdot 6})^{12} = (10^{12})^{12} = 10^{144}$$

**Second term:**

$$\begin{aligned} (10^{-8} \times (0.1)^6)^{10} &= (10^{-8} \times (10^{-1})^6)^{10} = (10^{-8} \times 10^{-6})^{10} = \\ &= (10^{-8-6})^{10} = (10^{-14})^{10} = 10^{-140} \end{aligned}$$

**Overall expression:**

$$\begin{array}{ccccccc} 10^{144} & \times & 10^{-140} & = & 10^{144-140} & = & 10^4 = 10000 \\ \uparrow & & \uparrow & & & & \\ \text{First term} & & \text{Second term} & & & & \text{as required.} \end{array}$$

## 1.2 Exponentials

A common mathematical application for powers are **polynomials** such as  $x^7 + 3x^2$ . The emphasis lies on the base being a variable, here  $x$ , while the power/exponent is fixed, here 7 respectively 2. We will see more on this in later chapters.

However for **exponentials** we view the base as fixed and the exponent as the variable. Fortunately calculations with exponentials follow the rules of powers discussed. Example:  $e^x$

**Worked Example 1.7.** Express  $\frac{1}{3^{-4x}}$  in the form  $a^x$ , for a suitable number  $a$ .

**Solution :**

$$\begin{aligned} \text{From Rule 6: } \frac{1}{3^{-4x}} &= 3^{4x} = \underset{\substack{\uparrow \\ \text{Rule 4}}}{(3^4)^x} = (\underbrace{3 \cdot 3}_9 \cdot \underbrace{3 \cdot 3}_9)^x \\ &= 81^x \\ &\text{the number } a \text{ is } 81. \end{aligned}$$

### Applications of exponentials:

In **biology and ecology** exponentials are used to model unconstrained growth of bacteria. After a fixed unit time, say 1 hour, a cell splits into two new cells. Hence every hour the total number of bacteria doubles. Thus after  $T$  hours there are  $2^T$ -times as many bacteria.

In **computer science** Moore's law (See, e.g. on wikipedia.org under "Moore's law") is the observation that the density of transistors in an integrated circuit doubles roughly every two years. Hence in  $y$  years the transistor density increases by  $2^{(y/2)}$ , and in particular it grows by a factor of  $\sqrt{2}$  every year.

## 1.3 Logarithms

In the example on bacterial growth we might ask when are there 8 times as many bacteria as at the start. In other words we are interested in the time  $T$  such that  $2^T = P$ , where  $P = 8$ .

**Worked Example 1.8.** i) Find  $T$  by recognition such that  $2^T = P$ , where  $P = 8, 64, 1024$ .

**Solution :**

$$\begin{aligned} &\text{Find } T, \text{ where} \\ \text{i) } 2^T &= 8 = 2^3 \quad \longrightarrow \quad T = 3 \\ \text{ii) } 2^T &= 64 = 2^6 \quad \longrightarrow \quad T = 6 \\ \text{iii) } 2^T &= 1024 = 2^{10} \quad \longrightarrow \quad T = 10 \end{aligned}$$

**Remark:** The underlying approach of solving exponential problems by *recognition* lies the important idea of *comparing of exponents*.

Suppose in the following  $\Delta$  and  $\square$  are expressions in terms of quantities that we are interested in. Suppose we are asked to solve the equation

$$a^\Delta = a^\square \quad (1.11)$$

Then the method of *comparing of exponents* states that

$$\Delta = \square \quad (1.12)$$

must be true, i.e. the exponents must agree.

Thus we are able to solve our original equation (1.11) by solving the equation (1.12) instead.

**Worked Example 1.9.** Find  $x$  by recognition such that

a)  $7^x = 49$ ;   b)  $3^x = 81$ ;   c)  $10^x = 100,00$ ;   d)  $5^x = 0.2$ ;   e)  $2^x = -8$

**Solution :** a)  $49 = 7 \times 7 = 7^2$  so  $7^x = 7^2$  gives  $x = 2$

$$b) 3^x = 81 = \underbrace{3 \cdot 3}_9 \cdot \underbrace{3 \cdot 3}_9 = 3^4 \implies x = 4$$

$$c) 10^x = 10000 = \underbrace{10 \cdot 10 \cdot 10 \cdot 10}_{(4 \text{ zeros})} = 10^4 \implies x = 4$$

$$d) 5^x = 0.2 = \frac{2}{10} = \frac{1}{5} = 5^{-1} \implies x = -1$$

$$e) 2^x = -8 \quad \text{has no solution}$$

**Remark:** As we have just shown there are problems that we can solve without a calculator just using recognition. We can equally argue that some problems can not have a solution.

The main question that we are trying to answer above is:

*Given real numbers  $a$  and  $c$ , to which power, say  $x$ , do we need to raise  $a$  to get  $c$ ?*

*In other words which number  $x$  solves the equation  $a^x = c$ ?*

Above we solved this problem by recognition, however most times that is not possible for us. In order to invert the operation “raising  $a$  to the power of” we apply the operation **logarithm to base  $a$**  which is denoted by  $\log_a$ .

### Logarithms to the base $a$

The equation  $a^x = c$  is equivalent to the equation  $x = \log_a c$ .

**Worked Example 1.10.** Express the following exponential equations in terms of logarithms

a)  $7^x = 49$ ; b)  $3^x = 81$ ; c)  $10^x = 100,00$ ; d)  $5^x = 0.2$ ; e)  $2^x = -8$

**Solution :**

a)  $7^x = 49 \Rightarrow x = \log_7 49$

b)  $3^x = 81 \Rightarrow x = \log_3 81$

c)  $10^x = 10000 \Rightarrow x = \log_{10} 10000 = \log 10000$

d)  $5^x = 0.2 \Rightarrow x = \log_5 0.2$

e)  $2^x = -8 \Rightarrow x = \log_2 (-8)$

but logs are defined only on positive numbers so  $\log_2 (-8)$  is undefined.

The rules for powers and exponentials can be translated directly into logarithms:

#### Rules for logarithms

**Rule 1:**  $\log_a a = 1,$  (1.13)

**Rule 2:**  $\log_a 1 = 0,$  (1.14)

**Rule 3:**  $\log_a(xy) = \log_a x + \log_a y,$  (1.15)

**Rule 4:**  $\log_a(x^y) = y \log_a x,$  (1.16)

**Rule 5:**  $\log_a \frac{1}{a} = -1$  (1.17)

**Rule 6:**  $\log_a \frac{1}{a^n} = -n$  (1.18)

**Rule 7:**  $\log_a \sqrt[n]{a} = \frac{1}{n}$  ( $n \in \mathbb{N}$ ) (1.19)

**Rule 8:**  $\log_a \sqrt[n]{a^m} = \frac{m}{n}$  ( $n \in \mathbb{N}$ ) (1.20)

The most common logarithms have the base  $a = 10$  and  $a = e = 2.71828 \dots$ . In order to solve logarithms to another base  $a$  we can change the base using the following identity. Suppose we want to express  $\log_b$  with base  $b$  in terms of  $\log_a$  with base  $a$

#### Rule of change of base for logarithms

$$\log_b c = \frac{\log_a c}{\log_a b}$$



**Remarks:**

1. Both log terms on the right hand side use base  $a$ .
2. On both sides  $c$  is in the higher position and  $b$  is in the lower position.

In order to remember and understand the rule it helps to look at its derivation:

**Worked Example 1.11.** Derive the rule for the change of base.

[non - examinable, but interesting!]

Take  $x = \log_b c$  so  $b^x = c$

We want to change the base from  $b$  to  $a$ , i.e.  
find  $a, y$  such that:  $b^x = a^y = c$  so  $y = \log_a c$ .

Also we note that  $b = a^{\log_a b}$

so  $b^x = (a^{\log_a b})^x = a^{x \log_a b}$  compare this to  $a^y$   
(rules for powers)

we see that  $y = x \log_a b$

But  $y = \log_a c$

So  $x \cdot \log_a b = \log_a c \Rightarrow x = \frac{\log_a c}{\log_a b} = \log_b c$

**Worked Example 1.12.** Solve  $4^x = 64$  using base  $a = 4$ ,  $a = 2$  and  $a = e$ .

**Solution :**

1. base  $a = 4$  :  $4^x = 64 = 4^3$  by recognition

or  $x = \log_4 64 = \log_4 (4^3) = 3 \underbrace{\log_4 4}_{= 1} = 3$

2. base  $a = 2$

$$x = \log_4 64 = \frac{\log_2 64}{\log_2 4} = \frac{6}{2} = 3$$

3. base  $e$

$$x = \log_4 64 = \frac{\ln 64}{\ln 4} = 3$$

**Worked Example 1.13.** Guesstimate the solution of  $7^x = 50$  then use a calculator.

**Solution :**

$$49 = 7^2$$

$$343 = 7^3$$

$$2 < x < 3$$

$x$  - should be very close to 2

Guess :  $x = \frac{2.05}{2.01}$

$$7^x = 50 \rightarrow x = \log_7 50 = \frac{\ln 50}{\ln 7} = 2.0104 \text{ (4dp)}$$

**Worked Example 1.14.** Simplify  $\log_3 x + \log_9 \left(\frac{1}{x}\right) + \log_{\sqrt{3}} x$  using a base  $a = 3$ .

**Solution :**

$$\log_3 x + \log_9 \left(\frac{1}{x}\right) + \log_{\sqrt{3}} x$$

need to change the base to  $a = 3$

$$\text{so } \log_9 \left(\frac{1}{x}\right) = \frac{\log_3 \frac{1}{x}}{\log_3 9} = \frac{\log_3 \frac{1}{x}}{\log_3 3^2} = \frac{\log_3 \frac{1}{x}}{2 \log_3 3}$$

$$\text{and } \log_{\sqrt{3}} x = \frac{\log_3 x}{\log_3 \sqrt{3}} = \frac{\log_3 x}{\log_3 3^{1/2}} = \frac{\log_3 x}{1/2}$$

so all together we get:

$$\begin{aligned} \log_3 x + \frac{\log_3 \frac{1}{x}}{2} + \frac{\log_3 x}{1/2} &= \log_3 x + \frac{1}{2} \log_3 (x^{-1}) + 2 \log_3 x \\ &= \log_3 x - \frac{1}{2} \log_3 x + 2 \log_3 x = \left(1 - \frac{1}{2} + 2\right) \log_3 x \\ &= \frac{5}{2} \log_3 x = 2 \frac{1}{2} \log_3 x \end{aligned}$$

## 1.4 Solving equations with exponentials and logarithms

**Worked Example 1.15.** Given that  $\ln a = 2$  solve  $a^x = e^6$  for  $x$ .

**Solution :**

$$\begin{aligned}
 a^x &= e^6 && \text{Apply } \ln \text{ to both sides} \\
 \ln a^x &= \ln e^6 \\
 x \ln a &= 6 \underbrace{\ln e}_1 && \text{because } \ln e = \log_e e \\
 \text{so } x \ln a &= 6 \\
 \text{but } \ln a &= 2 \quad \text{so} && x \cdot 2 = 6 \rightarrow x = 3
 \end{aligned}$$

**Worked Example 1.16.** Find  $x$  such that  $\log_4(2^{9x+4}) = 4$ .

**Solution :**

1. Use the rules of logarithms

$$\begin{aligned}
 \log_4(2^{9x+4}) &= 4 \\
 \text{rule 4: } (9x+4) \log_4 2 &= 4 \\
 \text{rule 7: } (9x+4) \cdot \frac{1}{2} &= 4 \rightarrow 9x+4 = 8 \\
 9x &= 4 \\
 x &= \frac{4}{9}
 \end{aligned}$$

2. Change the base to  $a = 2$

$$\text{LHS} = \log_4(2^{9x+4}) = \frac{\log_2 2^{9x+4}}{\log_2 4} = \frac{9x+4}{2} = \frac{1}{2}(9x+4)$$

so  $\frac{1}{2}(9x+4) = 4$  and we proceed as above.

$$x = \frac{4}{9}$$

We will consider two types of equations with exponentials:

**Type 1)** Equations containing no sums on either side

**Method:**

1. Change all terms to a suitable common base, typically the smallest whole number.
2. Combine the terms on either side into a single power.
3. Compare the exponents, respectively the logarithms with respect to the common base.
4. Solve the resulting equation in  $x$ .
5. Check your solution solves the original equation.

**Worked Example 1.17.** Solve  $2^{2x} \times 8^{x+1} = 4^{3x}$  for  $x$ .

Check your solution does indeed solve the equation.

**Solution :**

$$\begin{aligned}
 1. \quad & 2^{2x} \times (2^3)^{x+1} = (2^2)^{3x} \quad \rightarrow \quad 2^{2x} \times 2^{3(x+1)} = 2^{6x} \\
 2. \quad & 2^{2x+3x+3} = 2^{6x} \quad \rightarrow \quad 2^{5x+3} = 2^{6x} \\
 3. \quad & 5x+3 = 6x \quad \rightarrow \quad 4. \quad x=3 \\
 5. \quad & 2^{2 \cdot 3} \times 8^{3+1} \stackrel{2}{=} 4^{3 \cdot 3} = 4^9 \\
 & 2^6 \cdot 8^4 = 4^3 \cdot 4^4 \cdot 2^4 = 4^{3+4+2} = 4^9 \quad \checkmark
 \end{aligned}$$

**Worked Example 1.18.** Show  $2^{3x} \times 8^{x+1} = 4^{3x}$  has no solution for  $x$ .

**Solution :**

$$\begin{aligned}
 1. \quad & 2^{3x} \times (2^3)^{x+1} = (2^2)^{3x} \\
 2. \quad & 2^{3x+3x+3} = 2^{6x} \\
 3. \quad & 6x+3 = 6x \quad \therefore \Rightarrow 3 \neq 0 \\
 & \text{no solution for } x
 \end{aligned}$$

**Type 2)** Equations containing sums of exponentials on one side or both

**Method:**

1. Change all terms to a suitable common base  $a$ , typically the smallest whole number.
2. Use the substitution  $u = a^x$ .
3. Solve the resulting equation, typically a polynomial in  $u$ .
4. List the solutions  $u_i$ .
5. If  $u_i > 0$  then  $x_i = \log_a(u_i)$ , while  $u_i \leq 0$  gives no solution for  $x$ .
6. Check your solution solves the original equation.

**Worked Example 1.19.** Solve  $e^{2x} + e^x = 2$  for  $x$ . Check your final answer.

**Solution :**

1. ✓

2.  $u = e^x$  ,  $u^2 = e^{2x}$

3.  $u^2 + u = 2 \rightarrow u^2 + u - 2 = 0$   
 $(u - 1)(u + 2) = 0$

4.  $u_1 = 1$  ,  $u_2 = -2$

5.  $u_1 > 0 \rightarrow x_1 = \log_e 1 = \ln 1 = 0$   
 $u_2 < 0 \rightarrow$  no solution for  $x$

6.  $x = 0$

$e^{2 \cdot 0} + e^0 = 1 + 1 = 2 \checkmark$

**Worked Example 1.20.** Solve  $2^{2x} + 8^{x+1} = 4^{2x}$  for  $x$ .

**Solution :**

$$1. \quad 2^{2x} + (2^3)^{x+1} = (2^2)^{2x} \quad \text{changing all terms to base 2}$$

$$2^{2x} + 2^{3(x+1)} = 2^{4x}$$

$$2^{2x} + 2^{3x} \times 2^3 = 2^{4x}$$

$$2^{2x} + 8 \times 2^{3x} = 2^{4x}$$

$$2. \quad \text{Let } u = 2^x \rightarrow x = \log_2 u$$

$$\text{so } u^2 + 8u^3 = u^4$$

$$u^4 - 8u^3 - u^2 = 0 \quad \leftarrow \text{polynomial equation}$$

$$3. \quad u^2(u^2 - 8u - 1) = 0$$

$$\downarrow$$

$$u^2 = 0$$

or

$$u_{\pm} = \frac{8 \pm \sqrt{68}}{2} = 4 \pm \frac{\sqrt{68}}{2}$$

$$\downarrow$$

$$4. \quad u = 0$$

$$u_+ = 8.123, \quad u_- = -0.123$$

$$5. \quad u = 0 \rightarrow \text{no solution for } x$$

$$u_- = -0.123 < 0 \rightarrow \text{no solution for } x$$

$$u_+ = 8.123 \rightarrow x = \log_2 u_+ =$$

$$\text{EXACT FORM} = \log_2 \left( 4 \pm \frac{\sqrt{68}}{2} \right) =$$

16

$$= 3.02$$

APPROXIMATE

FORM  $\rightarrow$

6. Check

**Worked Example 1.21.** Solve the following system of simultaneous equations.

$$2 \ln x = \ln y + \ln 3 \quad (1)$$

$$e^x e^y = e. \quad (2)$$

What restrictions are there on  $x$  and  $y$ ? Explain.

**Solution:** Restrictions on  $x, y$ ?  
 $x > 0$  and  $y > 0$

$\ln(1)$   $\ln x^2 = \ln(y \cdot 3)$   
 compare expressions under the logs  
 ("recognition method")  
 $x^2 = 3y$

$\ln(2)$   $e^{x+y} = e^1 \rightarrow x+y=1$

so new system:

$$(3) \quad x^2 = 3y$$

$$(4) \quad \begin{cases} x^2 = 3y \\ x+y=1 \end{cases} \rightarrow y = 1-x \text{ sub into (3)}$$

$$x^2 = 3(1-x) = 3-3x$$

$$\Rightarrow x^2 + 3x - 3 = 0$$

$$x_{\pm} = \frac{-3 \pm \sqrt{21}}{2}$$

$$x_+ = \frac{-3 + \sqrt{21}}{2} > 0$$

$$x_- = -\frac{3 - \sqrt{21}}{2} < 0 \leftarrow \text{not a solution}$$

$$\text{so } y = 1 - x_+ = 1 - \frac{-3 + \sqrt{21}}{2} = \frac{5 - \sqrt{21}}{2}$$

so the solution is:  $x = \frac{-3 + \sqrt{21}}{2}, y = \frac{5 - \sqrt{21}}{2}$

## 1.5 Two applications of logarithms in science – NON EXAMINABLE

### 1) The Carbon-14 dating method

This method measures the percentage of  $C^{14}$  isotopes left in bone samples to date them. It uses the fact that the carbon isotope  $C^{14}$  has a half life of 5730 years, in other words after 5730 years the amount of  $C^{14}$  in a sample has shrunk to 50% of the initial amount.

Let  $t$  be the age of the sample measured in years, while  $\tau$  measures the age in half lives (HL). Hence we have  $\tau = 5730 t$ . In particular:

$$\tau = 1\text{HL}, t = 5730\text{yrs: } 50\% \text{ of } C^{14} \text{ left.}$$

$$\tau = 2\text{HL}, t = 11460\text{yrs: } 25\% \text{ of } C^{14} \text{ left.}$$

etc.

The percentage  $P$  of  $C^{14}$  left after  $\tau$  HL is given by:

$$P = (0.5)^\tau.$$

Hence

$$\tau = \log_{0.5} P = \frac{\log P}{\log 0.5} = \frac{\ln P}{\ln 0.5}$$

**Worked Example 1.22.** A bone sample is found to have 60% of  $C^{14}$  left in it.

- Guesstimate the age of the sample.
- Calculate the exact age of the bone sample.

- 60% is  $\frac{4}{5}$  of the way from 100% down to 50%.

so guess  $\frac{4}{5}\text{HL} = 4584 \text{ yrs}$

- Solution :

$$\tau = \frac{\ln 0.6}{\ln 0.5} = 0.737 \quad (3\text{dp})$$

$$\tau = 0.737 \text{ HL} \Rightarrow t = 4222 \text{ yrs}$$



**2) pH-values and the Nernst equation**

The pH-value is related to the concentration of the positive hydrogen ions  $H^+$  present in a solution. If  $[H^+]$  denotes the concentration of  $H^+$  then the **pH-value** is defined by

$$pH = -\log[H^+].$$

The pH-value is typically measured using electrodes. Let  $E$  measure the electrode potential,  $R$  be the gas constant,  $T$  the temperature in kelvin and  $F$  the Faraday constant. Moreover let  $E_0$  measure the standard electrode potential. Then the **Nernst equation** relates the electrode potential  $E$  and  $[H^+]$  the concentration of hydrogen ions by

$$E = E_0 + \frac{RT}{2F} \ln([H^+]^2). \quad (1.21)$$

**Worked Example 1.23.** Rearrange the Nernst equation to get an expression for the pH-value in terms of  $E, E_0, R, T$  and  $F$

Observe :  $\ln([H^+]^2) = 2\ln([H^+])$

Now  $E = E_0 + \frac{RT}{2F} \cdot 2\ln([H^+])$

$$E - E_0 = \frac{RT}{F} \ln([H^+])$$

$$\frac{F}{RT} (E - E_0) = \ln([H^+])$$

$$pH = -\log[H^+] = -\frac{\ln[H^+]}{\ln 10} \rightarrow \ln[H^+] = -pH \cdot \ln 10$$

change of base

so  $\frac{F}{RT} (E - E_0) = -pH \cdot \ln 10$

then  $pH = -\frac{F}{RT \ln 10} (E - E_0) =$

$$= \frac{F (E_0 - E)}{RT \ln 10}$$

**Learning outcomes**

- Knowing and applying the rules for powers and exponentials.
- Simplifying expressions involving powers with the same basis.
- Simplifying expressions involving powers with different bases.
- Understanding the relationship between exponential and logarithmic expressions.
- Transforming exponential expressions into logarithmic ones, and vice versa.
- Knowing and applying the rules for logarithms.
- Knowing and applying the rules for the change of base for logarithms.
- Knowing and applying the rules for logarithms.
- Simplifying logarithmic expressions.
- Understanding and applying the methods for solving different kinds of problems involving exponentials or logarithms.