

New Assessment 1: Deadline Friday Week-8

Nech-8

17:00 on 15m

Nov

Group Theory

Week 2, Lecture 1, 2 & 3

Dr Lubna Shaheen

Assessment 1 available Monday 11th Nov 10:00 am

Table of Contents

- **1** Symmetries of geometric objects
- 2 Symmetric group
- **3** Cyclic groups
- Integer modulo n Group
- 5 The Quarternion group
- 6 Matrix group
 - The general linear group $GL_n(\mathbb{F})$

\$1, 6, 6², 6³ \$. 68, 6²8, 6

- Upper Triangular Matrix
- **7** Exams Style Questions



Example: The group of symmetries of a hexagon is known as the dihedral group, denoted as D_{12} consists of the symmetries that can map the hexagon onto itself. These symmetries include:

Rotational Symmetries: Rotation by 0°, 60°, 120°, 180°, 240°, 300°.

Reflective symmetries: There are 6 axes of symmetry for a regular hexagon, 3 axes passing through opposite vertices, 3 axes passing through midpiunts of opposite sides.



Examples of Groups $H' = \{1, 1, 8\}$ another cycli Subgroup $H = \{1, 6, 8\}, 6, 6, 5\}$ is a subgroup of T_2 $G_{1}: \Lambda^{3} \Lambda^{2} = \Lambda^{5} \in H \quad closed \qquad Q_{12}.$ $G_{12}: \quad Associatewith \qquad \Lambda(\mathbb{R}^{4} \circ \mathbb{E}^{5}) = \Lambda(\mathbb{R}^{9}) = \Lambda(\mathbb{R}^{6}, \mathbb{R}^{3}) = \Lambda(\mathbb{R}^{6}, \mathbb{R}^{3})$ $= \Lambda(\mathbb{R}^{6}, \mathbb{R}^{3})$ $: \quad \mathcal{R}^{\circ} = I \in H \qquad (\mathbb{R} \circ \mathbb{R}^{4}) \circ \Lambda^{5} = \mathbb{R}^{5} \cdot \mathbb{R}^{5} = \mathbb{R}^{10}$ $: \quad \mathcal{R}^{\circ} = I \in H \qquad (\mathbb{R} \circ \mathbb{R}^{4}) \circ \Lambda^{5} = \mathbb{R}^{5} \cdot \mathbb{R}^{5} = \mathbb{R}^{10}$ $: \quad \mathcal{R}^{\circ} = I \in H \qquad (\mathbb{R} \circ \mathbb{R}^{4}) \circ \Lambda^{5} = \mathbb{R}^{5} \cdot \mathbb{R}^{5} = \mathbb{R}^{10}$ $G_{s}: \mathcal{L}=I \in H$ $G_4: k \cdot k^5 = 1, k^2 \cdot k^4 = 1, k^3 \cdot k^3 = 1$ cyclic _Subgroup.

The symmetry group of a subset $P \subseteq R^n$ depends on the geometric properties of P. It consists of all isometries that map P onto itself, preserving the distances between points in *P*. For finite sets, this often corresponds to a finite group of discrete symmetries (such as rotations and reflections), while for infinite or more structured sets like circles or spheres, the symmetry group can be continuous and infinite.

For any subset $P \subseteq \mathbb{R}^n$, the symmetry group Sym(P) is a subgroup of the isometry group of \mathbb{R}^n , which consists of all distance-preserving transformations in \mathbb{R}^n .

Translations, Rotations, Reflections, Rotoreflections *Lansformations we have not consoluted yet*.

Examples of Symmetry Groups for Specific Subsets *P*:

We mostly consider regulat polygones in R² • Points in \mathbb{R}^n **2** Line Segment in \mathbb{R}^2 **3** Regular Polygon in \mathbb{R}^2 1 96 P consists of a single point, then the Symmetry group is the Trivial group. Because the only Thansformation that preserve P 2) The symmetry group include is & Taviva. Reflection across the perpendicular A buseeter Rigo Rotation ____ 180°

Examples of Groups $\xi_{1,k}$ $\Im \quad \mathcal{D}_{8}, \mathcal{D}_{10}, \mathcal{D}_{12}, - \zeta_{10}$ (D(2): onthog no Group Rotations: by any angle around the Reflection: center of the ciscle. Acsors any line through the Contre of the curde.

K= 21,2,3, ... n? **Examples of Groups** Symmetric group: Group of all permutations on n symbols Suppose X - a finite set. Observation: Let X be a set, consider $f: X \to X$ the binary operation \circ = f(g(m))f-investible functions $f: K \rightarrow X_{h(x)}$ $(f \circ g)(x) = f(g(x))$ f(x) = 2 $f(x) = 5 \quad h: x \to x$ is associative $f \circ (g \circ h) = (f \circ g) \circ h$. Proof: (fo(goh)(x)) = fo(goh)(x)= fo(g(h(n)))f(3) = 1 $=f(g(h(\alpha)))$ f(n) = m(fog)oh(x) = (fog)(h(x)) = f(g(h(a)) / Associative.

Symmetric group: Group of all permutations on *n* symbols

Sym(K)

X-set , Sym(X)= {the collection of one-to-one and onto function $f : X \to X$ }

The symmetric group of degree n is the symmetric group on the set $X = \{1, 2, 3, \dots, n\} \text{ will be denoted by } S_n. \quad f: \{1, 2, 3, 4, 5\} \longrightarrow \{1, 2, 3, 4, 5\}$ Claim $f = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 1 & 5 & 4 & 3 \end{pmatrix}$ f(123 & 45) = (215 & 43)

Claim: Sym(X) equipped with \circ is a group.

G1: Closure: if f,geSym(X), fogeSym(X) G4: governe G1: Associative: we just cheched f': X->Y f': X->Y if f(Y) = X is the inverse of G3: Identity: f(a) = x $(g_{o}f_{i}(x) = g(f_{i}(x)) = g_{i}(x) = f(g_{i}(x)) = g_{i}(x)$

Verification of group axioms Symmetric group: Group of all permutations on *n* symbols

Disjoint cylce notation Writing permutations in disjoint

The group operation in a symmetric group is function composition, denoted by the symbol \circ or simply by just a composition of the permutations.

$$f = (13)(2)(45) = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 3 & 2 & 1 & 5 & 4 \end{pmatrix}$$
$$g = (125)(34) = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 5 & 4 & 3 & 1 \end{pmatrix}$$

We II apply first g and then f.

$$fg = f \circ g = (124)(35) = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 2 & 4 & 5 & 1 & 3 \end{pmatrix}$$

Dusjoint cycle Notation is not unique Though.

Verification of group axioms

Symmetric group: Group of all permutations on *n* symbols Remarks:

- Symmetric groups on infinite sets behave quite differently from symmetric groups on finite sets.
- **2** The symmetric group on a set of n elements has order n!.
- **③** It is abelian if and only if n is less than or equal to 2.
- For n = 0 and n = 1 (the empty set and the singleton set), the symmetric groups are trivial (they have order 0! and 1!.

The symmetric group on a set of size n is the Galois group of the general polynomial of degree n and plays an important role in Galois theory.

Symmetric group

 $e = \begin{pmatrix} 1 & 2 \\ 1 & 2 \end{pmatrix} f = \begin{pmatrix} -1 & 2 \\ 2 & 1 \end{pmatrix} \times = \begin{cases} 2 & 1 \\ 2 & 1 \end{cases}$

S_2 , symmetric group of degree 2

This group consists of exactly two elements: the identity and the permutation swapping the two points. It is a cyclic group and is thus abelian.

S_3 , symmetric group of degree 3, $S_3 \cong D_6$

 S_3 is the first non-abelian symmetric group. This group is isomorphic to the dihedral group of order 6, (D_6) the group of reflection and rotation symmetries of an equilateral triangle, since these symmetries permute the three vertices of the triangle. Cycles of length two correspond to reflections, and cycles of length three are rotations.

$$f = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix} \qquad \begin{cases} 2 & 1 \\ 2 & 1 \end{pmatrix} \qquad f^{2} = f^{2} = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 2 \\ 1 & 2 \end{pmatrix} = e$$

fog=gof Symmetric group $A \qquad k = R_{12}, \\ S =$ S3=D6= 1, 2, 2, 28, 28, 8} $S_{3} = \begin{cases} \binom{12}{3} \binom{12}{3} \binom{123}{132} \binom{123}{213} \binom{123}{23} \binom{12}{23} \binom{12}{23} \binom{12}{23} \binom{12}{23} \binom{12}{23} \binom{12}{23} \binom{12}{23} \binom{12}{3} \binom{1$

Symmetric group

Write f in disjoint cycles. $f = \begin{pmatrix} 12 & 3 & 4 & 5 & 6 & 7 & 8 \\ 4 & 8 & 7 & 6 & 5 & 3 & 1 & 2 \end{pmatrix} \in S_8$

f = (146371)(28)

Cyclic group

The cycic group of order *n*. $C_n = \{1, z_n, z_n^2, \cdots, z_n^{n-1}\}$ where z_n is an element of order n. unit circle Example: $z_n = e^{2\pi i/n}$. k = i $Z_n = k e^{2\pi i/n}$ {1}+{4] = 0 Another visualisation is $f'_{1} + [2] = [3]$ $\mathbb{Z}/n\mathbb{Z} = \{0, 1, \cdots, n-1\}$ $\mathbb{Z}/5\mathbb{Z} = \{0, 1, 2, \cdots, 4\}$ Group under with addition. $\mathbf{+}$ In=e = 2 e 2 Ti b/n | k=0,1,2,...,n-1 $= \frac{2}{2} \frac{\eta}{1}, e^{2\pi i/\eta}, e^{\pi i/\eta},$ n=

Cyclic group

Cyclic Groups: The infinite cyclic group

te infinite cyclic group

$$\mathcal{C}_{\infty} = \{1, z, z^{2}, z^{3}, \dots z^{-1}, z^{-2}, z^{-3}, \dots\} \qquad \text{modelo n.}$$

where z is an element of infinite order.

Example: (i) $\mathbb{Z} = \{0, +1, -1, +2, -2, +3, -3, \dots\}$ (ii) G-group, $g \in \mathcal{G}(g)$ is a cyclic group.

Alaz is not group. under mult oplication modele n.

(i) 1, generate the entire Group 1+1=2 $n = 1 \neq 1 \neq \cdots \neq 1 (n - t e s).$

-1 can also generale the group. n=n.1Exercise for you to Think about. $-\eta = -\eta \cdot I$

Integer modulo n Group Groups of units:

Consider the multiplicative subgroup Integers module n,

$$\begin{array}{c} S \cdot 7 = 35 \mod od(n) \\ = 11 \\ = 11 \\ \text{(numbers from 1 to } n - 1, \text{ which are co-prime to } n \}. \\ \mathcal{U}_{12} = \{1, 5, 7, 11\} \xrightarrow{\simeq} V_{4} \\ \mathbb{Z}/12\mathbb{Z} = \frac{2}{5}(1, 2, 35, 4, 5, \cdots, 5, 11)^{2} \\ \text{This is essentially equal to } \mathcal{V}_{4}. \\ \end{array} \qquad \begin{array}{c} \times \frac{1}{5} \frac{5}{5} \frac{7}{11} \\ \frac{1}{5} \frac{5}{5} \frac{1}{11} \frac{11}{7} \\ \frac{7}{7} \frac{7}{11} \frac{1}{15} \frac{5}{11} \\ \frac{11}{11} \frac{7}{5} \frac{7}{11} \\ \frac{7}{11} \\ \frac{7}{11} \frac{7}{11} \\ \frac{7}{11} \frac{7}{11} \\ \frac{7}{11} \\ \frac{7}{11} \frac{7}{11} \\ \frac{$$

a mod (n) Integer modulo *n* Group f_a , fa: KIng -> KIng falas= a.x fa is investible to **Result**: U_n is a subgroup of Sym $(\mathbb{Z}/n\mathbb{Z})$. a is copsing G1: a, be Un a, b are coprime to n ton a.b mod(n) EUn faof6 is also a permutation which is co-prime toin. 62: U2: G13: Jolombily fi: Hnzz -> Hnzz frofa=fa faofic fa. G3: Every element have unverse in the Un.

The Quaternion group

 $Q_{g} \not\equiv \mathcal{Q}_{g}$

The quaternion group Q_8 (sometimes just denoted by Q) is a **non-abelian** group of order eight, isomorphic to the eight-element subset $\{1, i, j, k, -1, -i, -j, -k\}$ of the quaternions under multiplication.

Quaternion group multiplication table (simplified form)



i = h $i^2 = j^2 = k^2 = -1$, $j \cdot \bar{z} = -\lambda$ -1.i = -i, -1.i = -i, -1.k = -ki.j = k, k = i, k.i = j $i \cdot i = -k \cdot k \cdot i = -i \cdot i \cdot k = -i$ hi=j $i \cdot h = -j$ $j \cdot i = -h h$ k = i $j \cdot h = i$

The Quaternion group

$$1 \rightarrow \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, i \rightarrow \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$$
$$j \rightarrow \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, k \rightarrow \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix} - k = \begin{pmatrix} 0 & -\tilde{z} \\ -\tilde{z} & 0 \end{pmatrix}$$

 $-i = \begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix}$

Exercire: Check, if This list of Subgroups of Q8 is complete.

Matrix group: General linear group, $GL_n(R)$ or GL(n, R).

Definition

In mathematics, the general linear group of degree n is the set of $n \times n$ invertible matrices, together with the operation of ordinary matrix multiplication. This forms a group, because the product of two invertible matrices is again invertible, and the inverse of an invertible matrix is invertible, with the identity matrix as the identity element of the group.

Remark: A matrix is invertible iff its determinant is not 0.

 $GL_{n}(\mathbb{K})$

n=2

R. C, Fp = Z/PZ = 30,1,2,..., P-15

Matrix group: General linear group, $GL_n(\mathbb{R})$ or $GL(n, \mathbb{R})$.

Real numbers case: The general linear group $GL(n, \mathbb{R})$ over the field of real numbers is a real Lie group of dimension n^2 . To see this, note that the set of all $n \times n$ real matrices, $M_n(\mathbb{R})$. forms a real vector space of dimension n^2 . The subset $GL(n, \mathbb{R})$ consists of those matrices whose determinant is non-zero.

Complex numbers case: The general linear group over the field of complex numbers. GL(n. \mathbb{C}), is a complex group of complex dimension n^2 . As a real Lie group (through realification) it has dimension $2n^2$. The set of all real matrices forms a real Lie subgroup. These correspond to N = Examinable $GL(n, \mathbb{R}) < GL(n, \mathbb{C}) < GL(2n, \mathbb{R})$ Support of the second se the inclusions

which have real dimensions n^2 , $2n^2$, and $4n^2 = (2n)^2$.

Matrix group: General linear group, $GL_n(\mathbb{R})$ or $GL(n, \mathbb{R})$.

Finite field case: Let \mathbb{F} be a field, then the general linear group $GL_n(\mathbb{F})$ consists of all the invertible $n \times n$ matrices over \mathbb{F} , the operation key matrix multiplication.

Exercise: $\mathbb{F}_2 = \{0,1\}.$ $\mathbf{\mathscr{V}}_{GL_2(\mathbb{F}_2)} = \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \right\}$ Non-examinable Question. **Exercise**: Comput $|GL_{100}(\mathbb{F}_2)|$. ? $|GL_n(F_q)| = q^n (q^n - q^n)(q^n - q) \dots (q^n - q^n)$ $|GL_{10}(F_2)| = 2^{\prime 0} (2^{\prime 0} - 2^{\prime}) (2^{\prime 0} - 2) - (2^{\prime 0} - 2^{\prime 7})$

Matrix group: General linear group, $GL_n(\mathbb{R})$ or $GL(n, \mathbb{R})$. **Definition**: The speciall linear group $SL_n(\mathbb{P})$ consists of $n \times n$ matrices with determinant 1. **Definition**: The special linear group $SL_2(\mathbb{R})$ is the group of 2×2 real matrices with det(A)=1 $det(A')=\frac{1}{det(A)}=1$ determinant 1. $Subgroup_{SL_2(\mathbb{R})} = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} : a, b, c, d \in \mathbb{R}, \text{ and } \underline{ad - bc} = 1 \right\} \Rightarrow A^{-l} \in SL_2(\mathbb{R})$ Suppose for the equation is infinite. Suppose for the equation is a group under multiplication. $G_1: A = \begin{pmatrix} a_1 & b_1 \\ c_1 & d_1 \end{pmatrix} \quad a_1 d_1 - b_1 c_1 = 1 \quad AB = \begin{pmatrix} q_1 q_2 + b_1 c_2, q_1 b_2 + b_1 d_2 \\ c_1 & d_1 \end{pmatrix}$ $B = \begin{pmatrix} a_1 & b_2 \\ c_2 & d_2 \end{pmatrix} \quad a_2 d_2 - c_2 b_2 = 1 \quad (c_1 q_2 + b_1 c_2, c_1 b_2 + d_1 d_2)$ $|SL_2(\mathbb{R})|$ is infinite. ABGSLn(P) ()() - ()() = 1

Subgroups of $SL_2(\mathbb{R})$:

Upper traingular matrix $geometry = I = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, det (I) = 1$ ICŠLn(P)

Cyclic Subgroups: Cyclic groups generated by one element of $SL_2(\mathbb{R})$. If $A \in SL_2(\mathbb{R})$, then $\langle A \rangle = \{A^n | n \in \mathbb{Z}\}$ forms a cyclic subgroups.

Diagonal Subgroups: The diagonal matrices with determinant 1 form a subgroup of

 $SL_2(\mathbb{R})$. This subgroup is isomorphic to \mathbb{R}^* , the multiplicative group of non-zero real $\begin{array}{c} \begin{pmatrix} i \\ a \\ o \\ a \end{pmatrix} \cdot A = \mathcal{T} \\ = \left\{ \begin{pmatrix} a & 0 \\ 0 & \frac{1}{2} \end{pmatrix} : a \in \mathbb{R} \right\}$ numbers: $A \cdot D = T$ det (A) = 1 $A = \begin{pmatrix} a_{1} & o \\ o & - \end{pmatrix} \quad B = \begin{pmatrix} a_{2} & o \\ o & - \end{pmatrix}$ $AB = \begin{pmatrix} a_1 a_2 + 0, 0 + 0 \\ 0 + 0, 0 + 1 \end{pmatrix} = \begin{pmatrix} a_1 a_2 & 0 \\ 0 & 1 \\ 0 & 0 \end{pmatrix} \in H \quad A^{-1} \in SL_2(\mathbb{R})$

Upper traingular matrix $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \not\subset I \qquad I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$

Upper Triangular Subgroup (Borel Subgroup):

The set of upper triangular matrices with determinant 1 forms an important subgroup, called the Borel subgroup: $P_{-}/9_{2}/6_{21}$

$$A = \begin{pmatrix} a_{1} & b_{1} \\ o & \frac{1}{a_{1}} \end{pmatrix} \qquad B = \left\{ \begin{pmatrix} a^{*} & b \\ 0 & \frac{1}{a} \end{pmatrix} : a \in \mathbb{R}^{*}, b \in \mathbb{R} \right\}$$
Rotation Subgroup (SO(2))
The special group
SO(2) is a subgroup of SL₂(\mathbb{R}), consisting of all rotations in \mathbb{R}^{2} :
$$SO(2) = \left\{ \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} : \theta \in \mathbb{R} \right\}$$

$$B = \begin{pmatrix} 1 & -b \\ \cos \theta \end{pmatrix} = \begin{pmatrix} 1 & -b \\ 0 & 4 \end{pmatrix} = \begin{pmatrix} 1 & -b \\ 0 & 4 \end{pmatrix} A A A^{*} = f$$

Upper traingular matrix

1

Exercise: Consider the **Upper triangular matrix with 1 on the diagonals**. Check if this is subgroup of $SL_2(\mathbb{R})$.

$$U = \begin{cases} (1 \ b) \\ 0 \ 1 \end{cases}; \ b \in \mathbb{R} \end{cases}$$

$$A = \begin{pmatrix} \cos 0 & -\sin 0 \\ \sin 0 & \cos 0 \end{pmatrix} \quad B = \begin{pmatrix} \cos 0 & -\sin 0 \\ \sin 0 & \cos 0 \end{pmatrix}$$

$$B = \begin{pmatrix} \cos 0 & \cos 0 \\ \sin 0 & \cos 0 \\ \sin 0 & \cos 0 \\ \sin 0 & -\sin 0 \sin 0 \\ \cos 0 & \sin 0 \\ -\sin 0 & \sin 0 \\ \cos 0 & \sin 0 \\ -\sin 0 & \cos 0 \\ -\sin 0 & \sin 0 \\ -\sin 0 & \cos 0 \\ -\sin 0 & \cos$$

Upper traingular matrix 🛛 🗠

Definition: The upper triangular matrix has all the elements below the main diagonal as zero. Also, the matrix which has elements above the main diagonal as zero is called a lower triangular matrix, also, written in the form of;

$$I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \qquad U = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1r} & a_{1,r+1} & \cdots & a_{1n} \\ 0 & a_{22} & \cdots & a_{2r} & c_{2,r+1} & \cdots & a_{2n} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & a_{rr} & a_{r,r+1} & \cdots \\ 0 & 0 & \cdots & 0 & a_{r+1,r+1} & \cdots \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & 0 & \cdots & a_{m,n} \end{pmatrix} = \begin{pmatrix} cos(-0) & -sin(-0) \\ Sin(-0) & Sin(-0) \\ -Sin(0) & Cos(-0) \end{pmatrix} = \begin{pmatrix} cos(0) & -sin(-0) \\ -Sin(0) & Cos(-0) \\ -Sin(0) & Cos(-0) \end{pmatrix} = \begin{pmatrix} cos(0) & -sin(-0) \\ -Sin(0) & Cos(-0) \\ -Sin(0) & Cos(-0) \end{pmatrix} = \begin{pmatrix} cos(0) & -sin(-0) \\ -Sin(0) & Cos(-0) \\ -Sin(0) & Cos(-0) \end{pmatrix} = \begin{pmatrix} cos(0) & -sin(-0) \\ -Sin(0) & Cos(-0) \\ -Sin(0) & Cos(-0) \\ -Sin(0) & Cos(-0) \end{pmatrix} = \begin{pmatrix} cos(0) & -sin(-0) \\ -Sin(0) & Cos(-0) \\ -Sin(0) & Cos(-0)$$

Upper traingular matrix

Properties of Upper Triangular Matrix

- If we add two upper triangular matrices, it will result in an upper triangular matrix itself.
- If we multiply two upper triangular, it will result in an upper triangular matrix itself.
- The inverse of the upper triangular matrix remains upper triangular.
- The transpose of the upper triangular matrix is a lower triangular matrix, $U^T = L$.
- If we multiply any scalar quantity to an upper triangular matrix, then the matrix still remains as upper triangular.

Exam Year, 2024

Question 1

(a) For the following, either give an example or explain why no example can exist: (i) A group with at least four elements in which every element has order either 1 or 2. (ii) A group with at least four elements in which every element has order either 1 or 4. (iii) Two groups of order 24 which are not isomorphic to one another. (iv) Two countably infinite groups which are not isomorphic to each other. (b) Let $G = \{x \in \mathbb{R} : x \ge 0\}$ and define a binary operation \circ on G by $x \circ y := |x - y|$. Decide which of the four group axioms are satisfied by (G, \circ) and which are not. For each axiom, give a brief justification for your answer. (i) V_{4} , \mathcal{D}_{4} $(g^{2})^{2} = 4$ g^{2} has order 2. (ii) Not possible, $1g\gamma = \frac{2}{5}1, g, g^{2}, g^{3}$

- (c) Using Lagrange's theorem, or otherwise, show that if g is an element of a group G such that |G| = n, then g^n is the identity element of G.
- (d) Using the result of (c) above, show that if p is a prime number and n is an integer in the range $1 \le n \le p$, then $n^{p-1} \equiv 1 \mod p$. (Hint: consider the group \mathcal{U}_{p} .)
- (e) List all subgroups of the dihedral group \mathcal{D}_{10} and indicate briefly why your list is complete.

b)
$$G_{1}: \chi_{70}, \chi_{70}, |\chi_{0}y| = |\chi_{-}y| = |2-3| = 1 \neq 0$$
 closed $G_{14}: \chi_{0}\chi = |\chi_{-}x| = 0$
 $|\chi_{-}y| \neq 0 = \neq \chi y \in G$
 $|\chi_{-}x| = 0$
 $G_{2}: Check are asocializing G_{3}: \pi_{00} = |\chi_{-}0| = |\chi| = \chi$
 $element$
 $\chi_{70}, \chi_{70}, \chi_{70}$
 0 is the identify.
 $|\chi_{-}g| = \chi_{-}y = 0$
 $|(\chi_{0}y) = ||\chi_{-}y| = 0$
 $||\chi_{0}(y_{0}y)| = ||\chi_{0}(y_{-}y)| = ||\chi_{-}(y_{-}y)| ||\chi_{-}y| = 0$

Question 2: Classify all groups of order 4 up to isomorphism.

Sol:Let G be a group with order |G| = 4. Then, we know by Lagrange's theorem that non-identity elements of G can have orders 2 or 4.

- If G contains an element of order 4, then G is cyclic and therefore isomorphic to \mathbb{Z}_4 .
- If G does not contain an element of order 4, the only other possibility is that all 3 non-identity elements have order 2. If we let G = {e, b₁, b₂, b₃}, we consider the value of b₁b₂. If b₁b₂ = e, then b₁ = b₂, a contradiction. If b₁b₂ = b₁ or b₁b₂ = b₂, then we conclude one of b₁ and b₂ is the identity, again a contradiction. So, we must have b₁b₂ = b₃. Then, we define a mapping φ : G → Z₂ × Z₂:

$$egin{aligned} arphi &: m{e}
ightarrow (0,0) \ arphi &: m{b}_1
ightarrow (0,1) \ arphi &: m{b}_2
ightarrow (1,0) \end{aligned}$$

 $arphi:b_3
ightarrow$ (1, 1)

giving us an isomorphism from ${\it G}$ to $\mathbb{Z}_2\times\mathbb{Z}_2$.

Therefore, every group *G* of order 4 is isomorphic to either \mathbb{Z}_4 or $\mathbb{Z}_2 \times \mathbb{Z}_2$.

Question 3: Classify all groups of order 8 up to isomorphism. Non-Examinable

Exam Year, 2022

Question: Give examples of

(a) A group of order 24 which is not abelian. \mathcal{D}_{24} ?

(b) A group of infinite order which is not abelian.

(c) A pair of abelian groups of the same order which are not isomorphic to one another.

(d) A group G and a two subgroups H_1 , $H_2 \leq G$ such that $H_1 \cup H_2$ is npt a subgroup of G.

Exams Style Questions Exam Year, 2021

Question 1 [16 marks].

(a) Suppose G is a set with three elements a, b, c, with a binary operation given by the following table.

$$\begin{array}{c|cccc}
a & b & c \\
\hline
a & b & a & c \\
b & a & b & c \\
c & c & c & b
\end{array}$$

Which of the group axioms G1–G4 does G satisfy? Justify your answer.

(b) Now let

$$H = \left\{ \begin{pmatrix} a & b \\ -b & a \end{pmatrix} \middle| a, b \in \mathbb{R}, a^2 + b^2 \neq 0 \right\} \cup \left\{ \begin{pmatrix} c & d \\ d & -c \end{pmatrix} \middle| c, d \in \mathbb{R}, c^2 + d^2 \neq 0 \right\}.$$

Prove that *H* is a subgroup of $GL_2(\mathbb{R})$.

Now suppose *G* is a group. Recall that if $g \in G$, the **order** of *g* is the smallest positive integer *n* such that $g^n = 1$, or ∞ if no such *n* exists.

(c) Suppose f,g ∈ G satisfy gf = f⁻¹g and ord(g) = 4. What is ord(fg)? Justify your answer.

[5]

[6]

Exams Style Questions Exams Question 2022

Recall that $GL_n(\mathbb{R})$ denotes the group of invertible $n \times n$ matrices with real entries. Let O(n) denote the set

$$O(n) = \{A \in GL_n(\mathbb{R}) : A^T A = I\}$$

where *I* denotes the $n \times n$ identity matrix and A^T denotes the transpose of the matrix *A*. Show that O(n) is a subgroup of $GL_n(\mathbb{R})$.

 $\underbrace{G_{I}: A, B \in O(n): A^{T}A = I}_{B^{T}B = I} \qquad \underbrace{G_{3}: I \in O(N)}_{I^{T}I = I} \\
 (AB)^{T}(AB) = B^{T}A^{T}AB = B^{T}IB \\
 = I$

 $A^{T}A = I = > A^{-1} = A^{T}$

Gig: AEO(n) $(A^T)^T A^T = A \cdot A^T = I$ => ATE O(N)

QMplus Quiz 2

= 3 1,8 2 Romembel | SR= 8 S | = 31, 6.8 7 (NS)(1S) = = \$ 1,28 ~ K(x's)-S=e Attempt Quiz 2 at QMplus page (183=1 = { 1, 1 3 3 } What about = {1,898} Subgroup generated 12 S. 12 S= by 22,877 L. L.S. L. L.S h3. h3= 2. h. 8. 2. K. 8 R2. 15 - R. S = h h. S. h.h. S h. 1. J. S. S= C = Nº K S. N. S => (225=1 = R. E'8.8= e

Some Useful Notations

Throughout this course, we use the following notation.

- C_n denotes the cyclic group of order n.
- Klein group often symbolized by the letter V₄ or as K₄ = ℤ₄ × ℤ₄ denotes the group {1, a, b, c}, with group operation given by

$$a^2 = b^2 = c^2 = 1$$
, $ab = ba = c$, $ac = ca = b$, $bc = cb = a$.

• U_n is the set of integers between 0 and *n* which are prime to *n*, with the group operation being multiplication modulo *n*.

Some Useful Notations

• \mathcal{D}_{2n} is the group with 2n elements

1,
$$r, r^2, \ldots, r^{n-1}, s, rs, r^2s, \ldots, r^{n-1}s$$

The group operation is determined by the relations $r^n = s^2 = 1$ and $sr = r^{n-1}s$.

- S_n denotes the group of all permutations of $\{1, \ldots, n\}$, with the group operation being composition.
- $GL_n(\mathbb{R})$ is the group of $n \times n$ invertible matrices with entries in \mathbb{R} , with the group operation being matrix multiplication.

•
$$Q_8$$
 is the group $\{1, -1, i, -i, j, -j, k, -k\}$, in which
 $i^2 = j^2 = k^2 = -1$, $ij = k$, $jk = i$, $ki = j$, $ji = -k$, $kj = -i$, $ik = -j$.