

Main Examination period 2019

MTH6104/MTH6104P: Algebraic structures II

Duration: 2 hours

Apart from this page, you are not permitted to read the contents of this question paper until instructed to do so by an invigilator.

You should attempt ALL questions. Marks available are shown next to the questions.

Calculators are not permitted in this examination. The unauthorised use of a calculator constitutes an examination offence.

Complete all rough work in the answer book and cross through any work that is not to be assessed.

Possession of unauthorised material at any time when under examination conditions is an assessment offence and can lead to expulsion from QMUL. Check now to ensure you do not have any unauthorised notes, mobile phones, smartwatches or unauthorised electronic devices on your person. If you do, raise your hand and give them to an invigilator immediately.

It is also an offence to have any writing of any kind on your person, including on your body. If you are found to have hidden unauthorised material elsewhere, including toilets and cloakrooms, it will be treated as being found in your possession. Unauthorised material found on your mobile phone or other electronic device will be considered the same as being in possession of paper notes. A mobile phone that causes a disruption in the exam is also an assessment offence.

Exam papers must not be removed from the examination room.

Examiners: Matthew Fayers and Alex Fink

© Queen Mary University of London (2019)

In this paper, we use the following notation.

- C_n denotes the cyclic group of order *n*.
- U_n is the set of integers between 0 and *n* which are prime to *n*, with the group operation being multiplication modulo *n*.
- \mathcal{D}_{2n} is the group with 2n elements

1, r, r^2, \ldots, r^{n-1} , $s, rs, r^2s, \ldots, r^{n-1}s$.

The group operation is determined by the relations $r^n = s^2 = 1$ and $sr = r^{n-1}s$.

- *S_n* denotes the group of all permutations of {1,...,*n*} (with the group operation being composition).
- Q_8 is the group $\{1, -1, i, -i, j, -j, k, -k\}$, in which

$$i^{2} = j^{2} = k^{2} = -1$$
, $ij = k$, $jk = i$, $ki = j$, $ji = -k$, $kj = -i$, $ik = -j$.

Question 1. [20 marks]

(a) Give the definition of a **group**.

Suppose *G* is a group and $f, g \in G$. In the rest of this question you may use elementary rules for manipulating powers of elements.

(b) Give the definition of the set $\langle g \rangle$, and prove that it is a subgroup of <i>G</i> .	[6]
--	-----

- (c) In the case of the group U_{25} , find all the elements of $\langle 6 \rangle$.
- (d) Give the definition of the **order** of *g*.
- (e) Suppose $\operatorname{ord}(f) = 3$, $\operatorname{ord}(g) = 4$ and $gf = f^2g$. What is the order of fg? Justify your answer. [5]

Page 2

© Queen Mary University of London (2019)

[3]

[4]

[2]

Question 2. [18 marks]

Suppose *G* is a group, $H, N \leq G$ and $g \in G$.

(a) Give the definition of the right coset <i>Hg</i> .	[2]
(b) Find all the right cosets of the subgroup $\{1, 9, 31, 39\}$ in \mathcal{U}_{40} .	[4]
(c) Define what it means to say that N is normal in G .	[2]
(d) Now suppose <i>N</i> is a normal subgroup of <i>G</i> . Give the definition of the set <i>NH</i> , and prove that <i>NH</i> is a subgroup of <i>G</i> .	[6]
(e) Give an example of a group <i>G</i> with $N, H \leq G$ such that <i>NH</i> is not a subgroup of <i>G</i> . [Ye do not have to prove that <i>N</i> and <i>H</i> are subgroups, but you should show that <i>NH</i> is not a	<i>5u</i>
subgroup.]	[4]

Question 3. [13 marks]

(a)	Give the definition of a transposition in S_n .	[2]
(b)	Give the definition of the alternating group A_n .	[2]
(c)	Suppose $h \in S_n$. Explain how you can use the disjoint cycle notation for h to find the order of h and to find whether $h \in A_n$. [You do not need to prove anything.]	[4]
(d)	Find an element of order 12 in A_9 , and write this element as a product of 3-cycles. [You do not need to prove anything.]	[5]

Question 4. [20 marks]

Suppose *G* and *H* are groups.

(a) Give the definition	of a homomorphism from <i>G</i> to <i>H</i> .	[2]
(b) Does there exist a h $\phi(j) = (4 \ 3 \ 2 \ 1)$? Just	homomorphism $\phi: \mathcal{Q}_8 \to \mathcal{S}_4$ such that $\phi(i) = (1\ 2\ 3\ 4)$ and stify your answer.	[5]
(c) Suppose $\phi : G \to H$ of ϕ .	<i>I</i> is a homomorphism. Give the definition of the image and kernel	[4]
(d) Give a precise state	ment of the First Isomorphism Theorem.	[3]
(e) Use the First Isomorean such that $C_{15}/K \cong C_{15}$	rphism Theorem to show that there is a normal subgroup K of \mathcal{C}_{15} \mathcal{C}_5 .	[6]

MTH6104/MTH6104P (2019)

MTH6104/MTH6104P (2019)

[6]


[4]

Page 4

Question 5. [19 marks]

(a) Suppose <i>G</i> is a group and <i>X</i> is a set. Give the definition of an action of <i>G</i> on <i>X</i> .	[3]
(b) Given an example of a transitive action of Q_8 on Q_8 . [You do not need to prove anything, but you should say clearly how the action is defined.]	[3]
(c) Suppose π is an action of <i>G</i> on <i>X</i> , and $x \in X$. Give the definition of the orbit containing <i>x</i> and the stabiliser of <i>x</i> .	[4]
(d) Give a precise statement of the Orbit-Stabiliser Theorem.	[3]
(a) Now let C be the summetry group of a triangular prism (with the triangular faces being	

(e) Now let *G* be the symmetry group of a triangular prism (with the triangular faces being equilateral):



What is |G|? Justify your answer.

Question 6. [10 marks]

Suppose *G* is a finite group and p is a prime number.

(a) Give the definition of a Sylow <i>p</i> -subgroup of <i>G</i> .	[2]
(b) Find a Sylow 2-subgroup of \mathcal{D}_{20} .	[4]

(c) Is this subgroup normal? Justify your answer.